

THE STRUCTURE OF A LOCAL EMBEDDING AND CHERN CLASSES OF WEIGHTED BLOW-UPS

ANCA M. MUSTĂŢĂ AND ANDREI MUSTĂŢĂ

ABSTRACT. For a local embedding of Deligne–Mumford stacks $g : Y \rightarrow X$, we find an étale cover of X by a (non-separated) Deligne–Mumford stack X' such that the fiber product $Y' = g(Y) \times_X X'$ is a finite union of étale covers Y'_i of Y , and such that the maps $Y' \rightarrow Y$ and $X' \rightarrow X$ are universally closed. Moreover, a natural set of weights on the substacks of X' allows the construction of a universally closed push-forward, and thus a comparison between the Chow rings of X' and X . In the case when X is the moduli space of stable maps or one of its intermediate spaces, the Chow ring of X' has been introduced and calculated in [MM1] as an extended Chow ring of a network. Here we identify the corresponding stack X' and its moduli problem in terms of stable maps with marked components. We apply the construction above to computing the Chern classes of a weighted blow-up along a regular local embedding via deformation to a weighted normal cone and localization. We apply these calculations to find the Chern classes of the stable map spaces.

INTRODUCTION

Local embeddings form an important class of morphisms of algebraic stacks. For instance, the morphisms from the components of the inertia stack of a Deligne–Mumford stack into the stack itself are in general local embeddings. As another fundamental example, the diagonal morphism of a stack is a local embedding and thus, the local study of this type of morphisms has led to a good definition of an intersection product on smooth Deligne–Mumford stacks by A. Vistoli [V], with a subsequent simplification by A. Kresch [K]. Their work relies on the existence, for any local embedding $Y \rightarrow X$, of an étale cover $V \rightarrow Y$, such that the composition $V \rightarrow X$ can be factored into a closed embedding $V \rightarrow U$ followed by an étale cover $U \rightarrow X$. Based on these covers, the normal cone of a local embedding ([V]), and a deformation of the ambient stack to the normal cone ([K]) can be constructed. However, these constructions are local in essence and as such, they fail to completely encode information on global invariants like Chern classes, or Chow ring structures, and cannot be directly employed in Riemann–Roch type theorems like the Riemann–Roch without denominators as formulated for closed embeddings of schemes.

In this paper we replace the local construction above by one more suited to the purposes just mentioned, at the expense of working with non-separated stacks rather than separated schemes. Theorem 1.9 highlights the existence, for every local embedding $g : Y \rightarrow X$, of an étale cover by a stack $X' \rightarrow X$, such that $Y' := g(Y) \times_X X'$, a finite union of étale covers Y'_i of Y , is embedded in X' , and such that the morphisms $Y' \rightarrow Y$ and $X' \rightarrow X$ are universally closed. Moreover, the morphism $X' \rightarrow X$ is an isomorphism

Date: February 6, 2009.

outside the image of Y . Thus the study of local embeddings can be reduced for practical purposes to that of closed embeddings of stacks.

While the map $p : X' \rightarrow X$ is forcefully not proper, it is universally closed. A weight function w on the set of substacks of X' is naturally attached to the map p , referring to the number of possible extensions of maps involved in the valuative criterion of properness. This "probabilistic weight" contributes to a good definition of push-forward p_* between the Chow rings of X' and X , an extension of the usual definition of proper push-forward to this type of universally closed maps. In effect, the Chow ring $A(X)$ is a subalgebra of $A(X')$ whose additive structure can be recovered from $A(X')$ via the universally closed push-forward p_* .

One advantage of our construction comes from the fact that for a suitable étale cover U of X , we take into account the entire pullback of that cover to the locally embedded Y . Locally, in the neighborhoods of some points in U , this pullback may consist of a number of intersecting components. Their intersections contribute essentially to the structure of the morphism $Y \rightarrow X$; in a first instance, to the associated flat stratification of X . For this reason we encode them in a network of morphisms of stacks, a stack version of the configuration schemes of [L]. In [MM1], we have defined the extended Chow ring of such a network. In Theorem 1.18 we prove that this extended Chow ring is isomorphic to $A(X')$. In particular, in the case of the moduli space of stable maps and the local embeddings of its boundary divisors, the extended Chow ring has been calculated in [MM1]. In Theorem 3.4 we now identify the corresponding stack $\overline{M}'_{0,m}(\mathbb{P}^n, d)$ and formulate its moduli problem in terms of stable maps with marked components.

Our approach is relevant, for instance, when considering a blow-up along a local embedding. Let us return to the initial picture of a commutative diagram

$$\begin{array}{ccc} V & \longrightarrow & U \\ \downarrow & & \downarrow \\ Y & \longrightarrow & X, \end{array}$$

where the top map is an embedding. Then simply taking the blow-up of U along V does not lead to an étale cover of a complete blow-up of X along Y , due to a break in symmetry at the level of relations. Indeed, in small enough neighborhoods around each point in U , one needs to consider blow-up along each of the components of $Y \times_X U$ before defining the étale cover for $\text{Bl}_Y X$.

Weighted blow-ups form a class of morphisms with a variety of applications. They come up, for example, when considering variation of GIT. As algebraic stacks provide a natural context for the study of weighted blow-ups, this class may be extended to weighted blow-ups along regular local embeddings. Moduli spaces of (weighted) curves and (weighted) stable maps are examples for which this type of morphisms comes up naturally.

As an application to the universally closed étale cover construction, blowing up Chern classes along a regular local embedding is reduced to the case of regular embeddings. The basic idea of this computation, like in the case of schemes, is to retrieve the Chern classes from their pullback to the exceptional divisor, for example via the

Riemann-Roch without denominators formula ([F]). However, when weights are considered, a less standard approach is necessary for the retrieval step. By a deformation to a "weighted normal cone", we reduce the problem to the case when both the blow-up locus and the exceptional divisor are fixed loci for $\mathbb{C}^*$ -actions making the blow-down morphism equivariant. The Atiyah-Bott localization theorem then means that the class of the exceptional divisor can be inverted, allowing us to retrieve a class on the blow-up from its pullback to the exceptional divisor.

The paper is organized as follows: In the first section we construct the universally closed morphism which turns a local embedding $Y \rightarrow X$ into an embedding. The first step is the case when the local embedding is étale on its image. The general case is reduced to this situation by flat stratification. A network of local embeddings depending on the étale structure of the strata is highlighted in section 1.2, and X' is constructed by induction on strata, such that all the local embeddings in the network are replaced by embeddings. In section 1.3, universally closed morphisms and push-forwards associated to probabilistic weights are defined. The existence of an isomorphism between the Chow rings of X' and of the correspondent network, as introduced in [MM1], is proven, and the relation between the Chow rings of X and X' is discussed. The example of the moduli space of stable maps is discussed in 1.4. The second section of the paper contains the calculation of Chern classes for weighted blow-ups. The third section is dedicated to the example of the stable map spaces and its intermediate weighted stable map spaces. The Appendix discusses the Euler sequence of a weighted projective bundle. Although the this sequence is most likely known, we could not find a proof in the literature, and thus decided to carefully trace the sequence through the groupoid presentation the weighted projective bundle.

Acknowledgements The initial motivation for this article came from a question of Jason Michael Starr. The first ideas were formulated during our stay at the Mathematical Sciences Research Institute in Berkeley, while the paper was finished during the second author's stay at the Max Planck Institute für Mathematik in Bonn. We are very grateful for the hospitality of both institutes. The authors were partially supported by a Science Foundation Ireland grant.

1. THE UNIVERSAL LIFT OF A LOCAL EMBEDDING

The stacks in this article are assumed to be algebraic in the sense of Deligne–Mumford, integral, Noetherian, and all morphisms considered between them are of finite type.

1.1. The lift of a local embedding étale on its image.

Definition 1.1. Following [V], we will call local embedding any representable unramified morphism of finite type of stacks. A regular local embedding is a local embedding which is also locally a complete intersection.

Given a local embedding $Y \rightarrow X$, by Lemma 1.19 in [V] there is a diagram

$$\begin{array}{ccc} V_1 & \xrightarrow{g_1} & U \\ \downarrow p_1 & & \downarrow p \\ Y & \xrightarrow{g} & X, \end{array}$$

the vertical morphisms being étale covers and g_1 being a closed embedding of schemes. Let V be the scheme representing the fiber product $Y \times_X U$, with the induced map $g : V \rightarrow U$, and the image of g denoted by W . As $V_1 \rightarrow Y$ and $V \rightarrow Y$ are étale, then so must be the induced morphism $i_1 : V_1 \rightarrow V$. Since $V_1 \rightarrow U$ is an embedding, then so must be $V_1 \rightarrow V$. We write $V = V_1 \cup V_2$, where V_2 is the closure in V of $V \setminus V_1$. In fact, V_1 and V_2 are disjoint, as the map $V = V_1 \cup V_2 \rightarrow Y$ is étale. Denote by p_i the restriction $p_i : V_i \rightarrow Y$, by W_i the image of V_i in U .

Furthermore, if the morphism g is étale on its image, then by the étale lifting property ([G] I, Proposition 8.1.), V_1 and U can be chosen such that

$$V_1 = g(Y) \times_X U,$$

and as such, there is an étale morphism $g : V \rightarrow V_1$ with the property that $g \circ i_1 = \text{id}_{V_1}$.

We recall the étale groupoid presentation $[R \rightrightarrows U]$ of the stack X , given by the two projection morphisms $R := U \times_X U \rightrightarrows U$, together with canonical morphisms e , m , and i . Here the identity e is the diagonal morphism $e : U \rightarrow U \times_X U$, the multiplication m is

$$m := \pi_{13} : U \times_X U \times_X U \cong (U \times_X U) \times_U (U \times_X U) \rightarrow U \times_X U,$$

and the inverse morphism $i : U \times_X U \rightarrow U \times_X U$ switches the two terms of the product.

Notation. Let

$$S_{ij} := \text{Im}(\phi_{ij} : V_i \times_Y V_j \rightarrow U \times_X U),$$

for the map ϕ_{ij} given as a composition

$$V_i \times_Y V_j \hookrightarrow V \times_Y V = V \times_Y (Y \times_X U) \cong V \times_X U \rightarrow U \times_X U.$$

We denote by R' the subset

$$R' := R \setminus (S_{12} \cup S_{21} \cup (S_{22} \setminus S_{11})) \cup \text{Im } e.$$

From now on let $g : Y \rightarrow X$ be a closed local embedding. Then S_{ij} are closed subschemes of R . In general $S_{22} \setminus S_{11}$ might not be closed, but it is so in the case when g is étale on its image, for dimension reasons. As U is an étale cover of X , then the diagonal $e : U \rightarrow U \times_X U$ is an open embedding, and thus, under the above condition on g , the subscheme R' of R is open.

Proposition 1.2. *Let $g : Y \rightarrow X$ be a closed local embedding, étale on its image. The restrictions $s_1, s_2 : R' \rightarrow U$ of the two projection morphisms $R \rightarrow U$, together with e , and with the restriction of m to $R' \times_U R'$ and of i to R' form the grupoid presentation of a Deligne-Mumford stack X' .*

Proof. First note that if $\{i, j\} = \{1, 2\}$, then $S_{12} \cap S_{11} = \emptyset$ and $S_{21} \cap S_{11} = \emptyset$. Indeed,

$$U \times_X U \hookleftarrow V_1 \times_X U \cong V_1 \times_Y (Y \times_X U) = V_1 \times_Y (V_1 \sqcup V_2) = (V_1 \times_Y V_1) \sqcup (V_1 \times_Y V_2).$$

We also note that R' may be written alternatively as a difference

$$R' := R \setminus (S_{12} \cup S_{21} \cup S_{22} \setminus (S_{11} \cup \text{Im } e_2)),$$

where e_2 is the composition of the diagonal $V_2 \rightarrow V_2 \times_Y V_2$ with the morphism $V_2 \times_Y V_2 \rightarrow U \times_X U$. Indeed, there is a sequence of consecutive Cartesian diagrams

$$\begin{array}{ccccc} V_2 & \hookrightarrow & V & \hookrightarrow & U \\ \downarrow & & \downarrow & & \downarrow \\ V_2 \times_Y V & \hookrightarrow & V \times_Y V & \hookrightarrow & U \times_X U, \end{array}$$

where the vertical maps are diagonal morphisms, with $V_2 \times_Y V_2 \subseteq V_2 \times_Y V$ and $V \times_Y V \cong V \times_X U$. This implies that $S_{22} \cap \text{Im } e = \text{Im } e_2$. Also, $S_{12} \cap \text{Im } e \subseteq S_{12} \cap S_{11} = \emptyset$, and symmetrically $S_{12} \cap \text{Im } e = S_{21} \cap \text{Im } e = \emptyset$.

As by construction R' is both symmetric and reflexive, the existence of a groupoid structure $[R' \rightrightarrows U]$ reduces to checking the closure of R' under multiplication. We note that for $k, l \in \{1, 2\}$, the preimage

$$\pi_{13}^{-1}(S_{kl}) \subseteq W \times_X U \times_X W \cong (W \times_X U) \times_U (U \times_X W).$$

But $W \times_X U$ is the image of $V \times_X U \cong \sqcup_{i,j \in \{0,1\}} V_i \times_Y V_j$, namely $\cup_{i,j \in \{0,1\}} S_{ij}$.

The injectivity of $g_1 : V_1 \rightarrow U$ directly implies $\pi_{13}(S_{11} \times_U S_{11}) \subseteq S_{11}$ while $\pi_{13}(S_{11} \times_U \text{Im } e_2) \subseteq S_{11} \cap S_{22}$ and $\pi_{13}(\text{Im } e_2 \times_U \text{Im } e_2) \subseteq \text{Im } e$. Thus the multiplication m is well defined on R' .

□

Remark 1.3. Elementary constructions similar to the above can be done directly in some cases when the local embedding $g : Y \rightarrow X$ is not étale on its image. A nice example is the case when an étale cover $V = V_1 \sqcup V_2$ of Y can be found such that both V_i are embedded in an étale cover U of X and $V \cong Y \times_X U$. In this case a natural choice for R' is $R' = R \setminus (S_{12} \cup S_{21})$ as $S_{11} \cap S_{12} = S_{11} \cap S_{21} = S_{22} \cap S_{12} = S_{22} \cap S_{21} = \emptyset$ and the multiplication map is thus well defined. In the general case of a local embedding though, an indirect construction X' will better reflect the structure of the map $Y \rightarrow X$. However, Proposition 1.2 will prove instrumental in that construction.

With the notations from the previous proposition, the following also holds.

Proposition 1.4. *There exist a natural embedding $Y \hookrightarrow X'$ and an étale map $X' \rightarrow X$ making the following diagram Cartesian*

$$\begin{array}{ccc} Y & \longrightarrow & X' \\ \downarrow & & \downarrow \\ g(Y) & \longrightarrow & X \end{array}.$$

Proof. The composition of étale morphisms $R' \hookrightarrow R \rightrightarrows U$ being étale, the natural morphism of groupoids

$$\begin{array}{ccc} R' & \longrightarrow & R \\ \Downarrow & & \Downarrow \\ U & \longrightarrow & U \end{array}$$

induces an étale morphism of stacks $X' \rightarrow X$. Next we check that $V_1 = Y \times_{X'} U$, namely that the diagram defining a groupoid morphism

$$\begin{array}{ccc} V_1 \times_Y V_1 & \longrightarrow & R' \\ \Downarrow & & \Downarrow \\ V_1 & \longrightarrow & U \end{array}$$

is Cartesian. Indeed,

$$V \times_Y V \cong V \times_X U \cong V \times_U (U \times_X U) = V \times_U R$$

and therefore

$$V \times_U R' \cong (V \times_Y V) \times_R R' \cong (V_1 \times_Y V_1) \sqcup \phi_{22}^{-1}(S_{11} \cup \text{Im } e_2),$$

and by restriction to V_1 ,

$$V_1 \times_U R' \cong V_1 \times_Y V_1.$$

This proves the existence of a natural embedding of Y in X' such that its composition with the étale map $X' \rightarrow X$ yields the local embedding of Y into X .

At the beginning of this section, the cover V_1 of Y was constructed as a fiber product $V_1 = g(Y) \times_X U$, and thus $V_1 \times_Y V_1 \cong V_1 \times_U R' \cong g(Y) \times_X R'$, forming an isomorphism of groupoids

$$[V_1 \times_Y V_1 \rightrightarrows V_1] \cong [g(Y) \times_X R' \rightrightarrows g(Y) \times_X U],$$

which induces an isomorphism $Y \cong g(Y) \times_X X'$. $\square$

Although the groupoid presentation of the stack X' depended on particular choices of covers for X and Y , the stack X' is uniquely defined by a universality property, which can be expressed in terms of moduli problems as follows

Theorem 1.5. *There is an equivalence of categories from the category of morphisms $Z \rightarrow X'$ defined on stacks Z of finite type to that of morphisms $Z \rightarrow X$ endowed with a section*

$$s : g(Y) \times_X Z \rightarrow Y \times_X Z$$

for the étale map $Y \times_X Z \rightarrow g(Y) \times_X Z$.

Proof. Indeed, at the level of objects, given a map $Z \rightarrow X'$ from a scheme of finite type Z into X' and its composition with the étale map $X' \rightarrow X$, there is an induced isomorphism

$$g(Y) \times_X Z \cong (g(Y) \times_X X') \times_{X'} Z \cong Y \times_{X'} Z,$$

and a natural map $Y \times_{X'} Z \rightarrow Y \times_X Z$ which, when composed with the étale map $g_Z : Y \times_X Z \rightarrow g(Y) \times_X Z \cong Y \times_{X'} Z$, yields the identity map.

Consider now a scheme of finite type Z and a morphisms $Z \rightarrow X$ represented by a morphism of groupoids

$$\begin{array}{ccc} R_Z & \longrightarrow & R \\ \Downarrow & & \Downarrow \\ U_Z & \longrightarrow & U \end{array}$$

Assume that there is a section $s : g(Y) \times_X Z \rightarrow Y \times_X Z$ of $g_Z : Y \times_X Z \rightarrow g(Y) \times_X Z$, represented by the groupoid map

$$\begin{array}{ccc} V_1 \times_U R_Z & \longrightarrow & V \times_U R_Z \\ \Downarrow & & \Downarrow \\ V_1 \times_U U_Z & \longrightarrow & V \times_U U_Z. \end{array}$$

Then the map $R_Z \rightarrow R$ factors through $R_Z \rightarrow R' \rightarrow R$. Indeed, as the restrictions of R and R' to $U \setminus \text{Im } V_1$ coincide, it is enough to show the factorization of groupoid maps

$$\begin{array}{ccccc} V_1 \times_U R_Z & \longrightarrow & V_1 \times_U R' & \longrightarrow & V_1 \times_U R \\ \Downarrow & & \Downarrow & & \Downarrow \\ V_1 \times_U U_Z & \longrightarrow & V_1 & \longrightarrow & V_1 \end{array}$$

The map $V_1 \times_U U_Z \rightarrow V_1$ is the composition of $V_1 \times_U U_Z \rightarrow V \times_U U_Z$ above with the natural maps $V \times_U U_Z \rightarrow V \rightarrow V_1$, while $V_1 \times_U R_Z \rightarrow V_1 \times_U R'$ is the composition

$$V_1 \times_U R_Z \rightarrow V \times_U R_Z \rightarrow V \times_U R \cong V \times_Y V \rightarrow V_1 \times_Y V \cong V \times_U R' \rightarrow V_1 \times_U R' \cong V_1 \times_Y V_1.$$

Together these maps induce the desired morphism $Z \rightarrow X'$. A direct check shows that the constructions above are functorial, and that the two functors constructed between the category $\text{Hom}(Z, X')$ and that of morphisms $Z \rightarrow X$ with the extra property specified in the hypothesis are inverse to each other. $\square$

Remark 1.6. Although the embedding $V_1 \hookrightarrow V$ does induce morphisms $V_1 \times_U U_Z \rightarrow V \times_U U_Z$ and $V_1 \times_U R_Z \rightarrow V \times_U R_Z$, together these induced morphisms do not in general form a morphisms of groupoids, as they are not necessarily compatible with the multiplication on R_Z . In particular, when $Z = X$, our construction does not imply the existence of a section for $Y \rightarrow g(Y)$, or for $X' \rightarrow X$.

1.2. The étale structure of a local embedding. Consider a local embedding of irreducible, proper, Noetherian stacks $g : Y \rightarrow X$. Let $\{S_i\}_i$ be the flattening stratification for the morphism $Y \rightarrow g(Y)$, and let $Y_n \hookrightarrow Y_{n-1} \hookrightarrow \dots \hookrightarrow Y_1 \hookrightarrow Y_0 = Y$ be a filtration of Y consisting of the closures $\overline{g^{-1}(S_i)}$ in Y , for all i as above. Thus the restrictions of g

$$g_n : Y_n \rightarrow g(Y_n), \quad g_{n-1}^o : Y_{n-1} \setminus Y_n \rightarrow g(Y_{n-1} \setminus Y_n), \quad \dots, \quad g_0^o : Y \setminus Y_1 \rightarrow g(Y \setminus Y_1)$$

are all étale.

Proposition 1.7. *There exist an étale cover U of X and closed subschemes $\{W_l\}_{l \in L}$ with intersections $W_I = \bigcap_{l \in I} W_l$ for $I \subset \mathcal{P}(L)$, such that*

$$\bigcup_{l \in L} W_l = g(Y) \times_X U, \quad \bigcup_{I \in \mathcal{P}_k} W_I = g(Y_k) \times_X U$$

for all $k = 0, \dots, n$, for some suitable choices of subsets $\mathcal{P}_k \subset \mathcal{P}(L)$, and such that for each I as above, there exist sets of étale covers $\{V_I^a\}_{a \in \{1, \dots, d_k\}}$ of Y_k with bijective morphisms $V_I^a \rightarrow W_I$ standing over $g_k : Y_k \rightarrow g(Y_k)$ satisfying

$$Y_k \times_X U = \bigcup_{I \in \mathcal{P}_k, a \in \{1, \dots, d_k\}} V_I^a.$$

Proof. The cover above can be constructed inductively on flattening strata. As a first step, an étale cover U_n of X can be chosen by successive application of étale lifting, such that $V_n := Y_n \times_X U = \sqcup_a V_n^a$, where each V_n^a is an étale cover of Y_n such that the lift $h_n : V_n \rightarrow U_n$ of g_n is an embedding on each V_n^a . If $Y_k \times_X U = \bigcup_{I \in \mathcal{P}_k, a \in \{1, \dots, d_k\}} V_I^a$ satisfying the required properties have been found for all $k > i$ for a suitable choice of cover U_{i+1} of X , then the covers $U_i \rightarrow X$ and $V_I^a \rightarrow Y_i$ with $I \in \mathcal{P}_i$ can be found by a local construction around the closed points in the image of $Y_i \times_X U_{i+1} \rightarrow U_{i+1}$. While for points in the image of $(Y_i \setminus Y_{i+1}) \times_X U_{i+1}$ étale lifting provides a suitable cover, for a closed point z of $\text{Im}(h_{i+1} : Y_{i+1} \times_X U_{i+1} \rightarrow U_{i+1})$, there is an open neighborhood $W_z \subset U_{i+1}$, an étale cover $W_z^1 \rightarrow W_z$, and an open subset V_z^1 in $Y_i \times_X W_z$ with an embedding $h_z^1 : V_z^1 \hookrightarrow W_z^1$. We take W_z^1 to be minimal, in the sense that

all its components are required to intersect the image of V_z^1 , otherwise they are simply discarded. The complement V_z of V_z^1 in $Y_i \times_X h_z^1(V_z^1)$ then is locally immersed in $h_z^1(V_z^1)$, and thus the previous steps can be repeated as needed. Moreover, the degree of the map $V_z \rightarrow h_z^1(V_z^1)$ at the preimages of z in $Y_{i+1} \times_X h_z^1(V_z^1)$ is smaller than that of the map $Y_i \times_X h_z^1(V_z^1) \rightarrow h_z^1(V_z^1)$. This guarantees the finiteness of the process. On the other hand, $Y_i \times_X (W_z^1 \setminus h_z^1(V_z^1))$ is also locally immersed in $W_z^1 \setminus h_z^1(V_z^1)$ and therefore the construction can be repeated in this direction as well. Finiteness of the process is guaranteed by the finiteness of the number of components of $g(Y_i \setminus Y_{i+1}) \times_X W_z$, while the fact that the stacks are Noetherian allows the construction above to be completed with the choice of a finite number of suitable closed points z as above. $\square$

Notation. For each index $I \in \mathcal{P}_k$ as above, consider the stack Y_I having V_I^a as an étale cover, with relations defined by increasing induction on k as follows. First let $Y_\emptyset := X$ with the given presentation $[R_\emptyset := U \times_X U \rightrightarrows U]$. When $I = \{i\} \in \mathcal{P}_1$, then $Y_i \cong Y$, having thus a groupoid presentation $R_i \rightrightarrows V_i^a$ where

$$R_i = V_i^a \times_Y V_i^a \cong W_i \times_X W_i \setminus \bigcup_{(j,b) \neq (i,a)} S_{ij}^{ab},$$

where $S_{ij}^{ab} = \text{Im} (V_i^a \times_Y V_j^b \rightarrow W_i \times_X W_j)$. Then for any $I \in \mathcal{P}_k$, define

$$R_I = \left(\prod_{i \in I} \right)_{R_\emptyset} R_i,$$

where $(\prod)_{R_\emptyset}$ denotes the fiber product over $R_\emptyset$. Thus for any K, J there is a natural isomorphism $R_{J \cup K} \cong R_J \cup_{R_{J \cap K}} R_K$. The groupoid structures $[R_i \rightrightarrows V_i^a \cong W_i]$ induce a groupoid scheme $[R_I \rightrightarrows V_I^a \cong W_I]$ over $W_I = (\prod_{i \in I})_U W_i$. Let Y_I denote the stack associated to this groupoid scheme. By induction we obtain that for a fixed index a ,

$$R_I = V_I^a \times_{Y_I} V_I^a \cong W_I \times_X W_I \setminus \bigcup_{(j,b) \neq (i,a); i \in I} S_{ij}^{ab}$$

Remark 1.8. Although the stacks Y_I are not fibered products of stacks Y_i with $i \in I$, they can be constructed intrinsically from a succession of fibered products, after removing the diagonal components. In particular, their definition above implies the existence of a closed embedding

$$Y_I \hookrightarrow \left(\prod_{i \in I} \right)_X Y_i,$$

and thus the stacks Y_I are proper, Noetherian.

Also from definition, $Y_{I \cup \{h\}} \times_{Y_I} V_i^a = \bigsqcup_{J \supset I, |J \setminus I|=1} V_J^a$, and thus an inductive calculation yields, for all $I \in \mathcal{P}_k$ and $J \in \mathcal{P}_l$ such that $I \supset J$

$$(1.1) \quad Y_I \times_{Y_J} V_J^a \cong \bigsqcup_{K_k \supset K_{k-1} \supset \dots \supset K_{l+1} \supset K_l = J} V_K^a,$$

including as many copies of V_K^a as there are chains $K_k \supset K_{k-1} \supset \dots \supset K_{l+1} \supset K_l = J$ with $K_l \in \mathcal{P}_l$. When $J = \emptyset$,

$$(1.2) \quad Y_I \times_X U \cong \bigsqcup_{a, K_k \supset K_{k-1} \supset \dots \supset K_1 \supset K_0 = \emptyset} V_K^a.$$

For any local embedding $Z \rightarrow T$ defined on an integral stack Z , let $[Z \rightarrow T]$ denote the degree of the fibre at the generic point of the image. For $K \supseteq J$ such that

$|K \setminus J| = k - j$, the following relations thus hold

$$\begin{aligned} [Y_K \rightarrow Y_J] &= \sum_{K=K_k \supset K_{k-1} \supset \dots \supset K_{j+1} \supset K_j=J} \prod_{i=1}^{j-k} [Y_{K_{i-1}} \rightarrow Y_{K_i}] \\ &= N(k, j) \prod_{i=1}^{j-k} [Y_{K_{i-1}} \rightarrow Y_{K_i}] \end{aligned}$$

for a fixed choice of chain, where $N(k, j)$ is the number of all chains $K = K_k \supset K_{k-1} \supset \dots \supset K_{j+1} \supset K_j = J$ as above. Thus if $K \in \mathcal{P}_k$, $J \in \mathcal{P}_j$ and $I \in \mathcal{P}_i$ satisfy $K \supset J \supset I$, then by a count of chains

$$(1.3) \quad \frac{[Y_K \rightarrow Y_J][Y_J \rightarrow Y_I]}{[Y_K \rightarrow Y_I]} = \frac{|\{K_k \in \mathcal{P}_k; K_k \supset I\}|}{|\{K_k \in \mathcal{P}_k; K_k \supset J\}| |\{K_j \in \mathcal{P}_j; K_j \supset I\}|}$$

Theorem 1.9. *Consider a network of local embeddings $\phi_J^I : Y_J \rightarrow Y_I$ for $I \subseteq J$, $I \in \mathcal{P}_i$ and $J \in \mathcal{P}_j$, associated to a local embedding $Y \rightarrow X$ by the construction above, where by convention $Y_\emptyset = X$. For each such morphism ϕ_J^I , there exists a unique embedding of stacks $\phi_J^I : Y_J' \hookrightarrow Y_I'$, together with étale morphisms $p_J : Y_J' \rightarrow Y_J$ and $p_I : Y_I' \rightarrow Y_I$ making the diagram*

$$\begin{array}{ccc} Y_J' & \xrightarrow{\phi_J^I} & Y_I' \\ p_J \downarrow & & \downarrow p_I \\ Y_J & \xrightarrow{\phi_J^I} & Y_I \end{array}$$

commutative, and such that

$$Y_J \times_{Y_I} Y_I' = \bigsqcup_{J' \in \mathcal{P}_j, J' \supseteq I} Y_{J'}.$$

Proof. We construct Y_I' in decreasing order of I , with $Y_I' = Y_I$ for I maximal. For $I \in \mathcal{P}_{n-1}$, the stack $N_I := \bigsqcup_{J \supset I} Y_J$ comes with an étale map $n_I : N_I \rightarrow Y_I$ and the étale cover

$$\bigsqcup_{J \supset I} W_J \cong n_I(N_I) \times_{Y_I} W_I.$$

Y_I' is constructed as in Proposition 1.2, such that it admits

- (1) a surjective étale morphism to $p_I : Y_I' \rightarrow Y_I$,
- (2) an embedding $N_I \hookrightarrow Y_I'$ whose composition with p_I is n_I ,
- (3) an étale cover W_I satisfying $Y_J' \times_{Y_I} W_I \cong W_J$, and
- (4) a groupoid presentation $R_I' \rightrightarrows W_I$ given by

$$R_I' \cong W_I \times_X W_I \setminus \bigcup_{(j,b) \neq (i,a); i \in I} S_{ij}^{ab} \setminus \bigcup_{(k,c) \neq (l,a); k, l \notin I} S_{lk}^{ac}.$$

Given any $k < n - 1$, $I \in \mathcal{P}_k$ and assuming the stack Y_J' with the above properties constructed for all $J \supset I$, the network stack N_I is defined out of all stacks $Y_{I \cup \{h\}}'$, by gluing each pair $Y_{I \cup \{h\}}'$ and $Y_{I \cup \{h'\}}'$ along $Y_{I \cup \{h, h'\}}'$ as in [AGV], Proposition A.1.1 and Corollary A.1.2. Accordingly, N_I admits a natural groupoid presentation $[\bigcup_{J \supset I} R_J' \rightrightarrows \bigcup_{J \supset I} W_J]$, where the unions are considered inside $U \times_X U$ and U , respectively. From here, the composition maps $Y_{I \cup \{h\}}' \rightarrow Y_{I \cup \{h\}} \rightarrow Y_I$ glue together to a morphism $N_I^I \rightarrow Y_I$ which

is étale on its image, and the construction can be repeated to obtain the stack Y'_I with the desired properties (1)–(4).

At the final step, the stack X' admits a groupoid presentation $[R' \rightrightarrows U]$, with

$$R' = U \times_X U \setminus \bigcup_{(i,a) \neq (j,b)} S_{ij}^{ab}.$$

□

1.3. Chow rings and universally closed pushforwards. The stacks Y'_I are in general non-separated. However, they do satisfy the existence part in the valuative criterion of properness.

Definition 1.10. A morphism of stacks $f : F \rightarrow G$ will be called universally closed if it is of finite type and, for any complete valuation ring R with field of fractions K and any commutative diagram

$$\begin{array}{ccc} \mathrm{Spec}(K) & \xrightarrow{u} & F \\ \downarrow & & \downarrow f \\ \mathrm{Spec}(R) & \longrightarrow & G, \end{array}$$

there exist finite extensions K' of K such that, for the integral closure R' of R in K' , the composition morphism $\mathrm{Spec}(K') \rightarrow F$ extends to $\mathrm{Spec}(R')$. Like in [V], we are mostly interested in stacks having coarse moduli schemes. Then by Proposition 2.6. and Definition 2.1 in [V], the image of such a morphism of stacks is defined and the definition above is a natural extension to stacks of the notion of universally closed morphism of schemes.

Definition 1.11. Consider a universally closed morphism of stacks $f : F \rightarrow G$. A probabilistic weight w of f is a map defined on the set of all integral substacks of F , with values in the interval $[0, 1]$, such that the weight of the generic point of F is 1 and, for any commutative diagram like in Definition 1.10,

$$w(D) = \sum_i w(P_i),$$

where D is the image in F of the unique point in $\mathrm{Spec}(K)$, and P_i are the images of the closed point in $\mathrm{Spec}(R')$ for all map extensions $\mathrm{Spec}(R') \rightarrow F$ as above.

Definition 1.12. Given a universally closed morphism of integral stacks $f : F \rightarrow G$ with probabilistic weight w , let

$$f_*[V] = w(V) \deg(V/W)[W]$$

for any closed integral substack V of F , where $W = f(V)$ and $\deg(V/W)$ is as in Definition 1.15, [V]. A homomorphism $f_* : Z_k(F) \rightarrow Z_k(G)$ is then defined by linear extension.

Proposition 1.13. *The homomorphism $f_* : Z_k(F) \rightarrow Z_k(G)$ induces a well defined universally closed push-forward homomorphism $f_* : A_k(F) \rightarrow A_k(G)$.*

Proof. Propositions 3.7 in [V] and Proposition 1.4, [F] deal with the proper push-forward in the cases of stacks and schemes, respectively. The difference for universally closed maps lies in the proof of the later, in the case when $\dim F = \dim G$. The case when f is finite is identical to Case 2 in Proposition 1.4, [F]. Following [F] closely, we take normalizations of the source and target, and the problem is thus reduced to the case of a universally closed morphism of normal varieties. Let W be a codimension one subvariety of G , let A be the local ring of W on G , and B the integral closure of the A in the field $k(F)$ of rational functions on F , such that B is a discrete valuation ring. Then by Definition 1.10, for each maximal ideal m_i of B there is a finite number of codimension one subvarieties V_i^l of F such that B dominates, and is therefore equal to, the local ring of each V_i^l in X . Then for any $r \in k(F)^*$,

$$\sum_i \text{ord}_{V_i^l}(r) \deg(V_i^l/W) = \text{ord}_W(N(r)),$$

where for each i a choice of the index l has been fixed, and $N(r)$ is the determinant of the $k(G)$ -linear endomorphism of $k(F)$ given by multiplication by r . Finally,

$$f_*[\text{div}(r)] = \sum_{V_i^l} \text{ord}_{V_i^l}(r) f_*[V_i^l] = \sum_{V_i^l} \text{ord}_{V_i^l}(r) w(V_i^l) \deg(V_i^l/W) [W].$$

For i fixed, $\text{ord}_{V_i^l}(r) \deg(V_i^l/W)$ is constant and $\sum_l w(V_i^l) = w(F) = 1$ from Definition 1.3. Thus

$$f_*[\text{div}(r)] = \sum_W \text{ord}_W(N(r)) [W]$$

like in the case of proper morphisms. This finishes the part of the proof specific to the universally closeness of f . $\square$

Universally closed push-forwards enjoy the usual properties of their proper relatives: for example, they commute with flat pullbacks, and the usual projection formula holds for f universally closed and flat.

Proposition 1.14. *Let $g : Y \rightarrow X$ be a local embedding and let X' be the corresponding étale, non-separated lift of X constructed in Theorem 1.9. Then the morphism $p : X' \rightarrow X$ is universally closed, and admits a probabilistic weight w .*

Proof. Let ξ_I denote the generic points of Y'_I , with $\xi_\emptyset$ the generic point of X' . With the notations from equation 1.3, define

$$w(\xi_I) := \frac{1}{|\mathcal{P}_k|[Y_I \rightarrow X]}$$

for any $I \in \mathcal{P}_k$. Then equation 1.3 becomes

$$(1.4) \quad w(\xi_I) = \sum_{J \in \mathcal{P}_k, J \supset I} [Y_J \rightarrow Y_I] w(\xi_J).$$

As $\{Y'_I \setminus (\bigcup_{J \supset I} Y'_J)\}_I$ forms a locally closed stratification of X' , the generic points of any integral substack D of X' will be found in exactly one of the above strata. We extend w to a function on all points of X' by identifying $w(D)$ with the weight of its associated stratum.

The universally closeness property will result during the proof that w is a probabilistic weight of $p : X' \rightarrow X$. Consider a complete discrete valuation ring R with field of fractions K , a commutative diagram

$$\begin{array}{ccc} \mathrm{Spec}(K) & \xrightarrow{u} & X' \\ \downarrow & & \downarrow p \\ \mathrm{Spec}(R) & \xrightarrow{v} & X, \end{array}$$

such that the image q_0 in X' of the generic point of $\mathrm{Spec}(R)$ lies in a stratum $Y'_I \setminus (\bigcup_{K \supset I} Y'_K)$, and the image q_1 in X of the closed point of $\mathrm{Spec}(R)$ lies in $\mathrm{Im}(Y_J \setminus \bigcup_{K \supset J} Y_K \rightarrow X)$ for some $J \supseteq I$. Since $(Y'_I \setminus (\bigcup_{K \supset I} Y'_K)) \cong (Y_I \setminus (\bigcup_{K \supset I} Y_K))$ and Y_I is proper, there is a unique extension $v' : \mathrm{Spec}(R') \rightarrow Y_I$ of a composition $\mathrm{Spec}(K') \rightarrow \mathrm{Spec}(K) \rightarrow X'$, with the notations of Definition 1.10. Let $q \in \mathrm{Im}(Y_J \rightarrow Y_I)$ be the lift of q_1 through this extension. Then through each point in the preimage of q in X' there is a unique lift in $Y'_I \hookrightarrow X'$ of the map v' . The generic point of each such lift has to be q_0 , because the map $p_I : Y'_I \rightarrow Y_I$ restricts to the above mentioned isomorphism $(Y'_I \setminus (\bigcup_{K \supset I} Y'_K)) \cong (Y_I \setminus (\bigcup_{K \supset I} Y_K))$. Let p_i be the images of the closed points of these lifts. They all have the same weight $w(\xi_J)$. Then by equation 1.4,

$$w(q_0) = \sum_i w(p_i),$$

which proves that w is a probabilistic weight for the universally closed morphism p . $\square$

Corollary 1.15. *For each $i \in \{0, \dots, n\}$ and $I \in \mathcal{P}_i$, there is a universally closed push-forward map $p_{[I*]} : A_k(Y'_I) \rightarrow A_k(Y_I)$, such that*

$$p_{[I*]} \circ p_{[I]}^* = d \cdot \mathrm{id} : A_k(Y_I) \rightarrow A_k(Y_I),$$

where d is the degree of the morphism $Y'_I \rightarrow Y_I$. In particular, the flat pullback $p_{[I]}^*$ is injective.

By convention, $X' = Y'_\emptyset$, and thus the Corollary shows how $A(X)$ can be regarded as a subalgebra of $A(X')$, and how classes in $A(X)$ can be recovered from $A(X')$ via push-forward.

Consider a local embedding $\phi : Y \rightarrow X$ with étale covers U for X and V for Y defined as in Section 1.2. Assume that all maps $\phi_I^J : Y_I \rightarrow Y_J$ are local regular embeddings. An extended Chow ring of the network $\{\phi_I^J : Y_I \rightarrow Y_J\}_{I,J}$, was introduced in Definition 3.6 of [MM1]. We recall this definition with a slight variation that does without the action of a symmetry group on $\mathcal{P}$.

Notation. Fix I and $I \cup \{h\} \in \mathcal{P}$. For any cycle $\alpha = [V] \in Z_l(\mathrm{Im}(Y_{I \cup \{h\}} \rightarrow Y_I))$, let $\alpha_h \in Z_l(Y_{I \cup \{h\}})$ be defined as follows:

$$\alpha_h = \frac{\sum_i [V_h^i]}{\deg((\phi_{Ih}^I)^{-1}(V)/V)}$$

where $\{V_h^i\}_i$ are the l -dimensional components of $(\phi_{Ih}^I)^{-1}(V)$.

Definition 1.16. The vector spaces $A_l(\{Y_I\}_{I \subseteq N}; \mathbb{Q})$ are defined by

$$A_l(\{Y_I\}_{I \subseteq N}; \mathbb{Q}) := \oplus_I Z^{l - \mathrm{codim}_Z Y_I}(Y_I) / \sim,$$

the sum taken after all $I \in \mathcal{P}$ with $\text{codim}_Z Y_I \leq l$. The equivalence relation $\sim$ is generated by rational equivalence together with relations of the type:

$$j_{Ih*}^I \alpha \sim \sum_{h'} \alpha_{h'},$$

for any cycle $\alpha \in Z_l(\text{Im}(Y_{I \cup \{h\}} \rightarrow Y_I))$.

The field of coefficients $\mathbb{Q}$ will be omitted in the notations throughout the rest of the text.

Definition 1.17. Multiplication is defined as

$$\alpha \cdot_r \beta := \bar{\phi}_{I \cup J}^{I*}(\alpha) \cdot \bar{\phi}_{I \cup J}^{J*}(\beta) \cdot c_{\text{top}}(\bar{\phi}_{I \cup J}^{I \cap J*} \mathcal{N}_{Y_{I \cap J}|Z})$$

in $A^*(Y_{I \cup J})$, for any two classes $\alpha \in A^*(Y_I)$ and $\beta \in A^*(Y_J)$. Here $\bar{\phi}_{I \cup J}^{I*}, \bar{\phi}_{I \cup J}^{J*}$ are the (generalized) Gysin homomorphisms, as defined in [V].

Theorem 1.18. *The following rings are isomorphic*

$$A(X') \cong A(\{Y_I\}_{I \subseteq N}).$$

Proof. Compositions of the flat pullbacks p_I^* with the push-forward of embeddings $\phi_{I*}^\emptyset$ add up to a morphism

$$\oplus_{I \subseteq N} A(Y_I) \rightarrow A(X').$$

which moreover factors through

$$F : A(\{Y_I\}_{I \subseteq N}) \rightarrow A(X').$$

Construct an inverse as follows. For each $J \subseteq N$, let U_J denote the complement in Y'_J of all the images of Y'_I , with $J \subset I$. (To define this complement one can work with supports of the corresponding coarse moduli schemes, but there is a canonical stack structure on U_I). Note that $U_J = Y_J \setminus (\bigcup_{I \subset J} \text{Im}(Y_I \rightarrow Y_J))$ as well. Working with the commutative diagram of open/closed exact sequences

$$\begin{array}{ccccccc} \oplus_{k \notin J} A(Y_{J \cup \{k\}}) & \longrightarrow & A(Y_J) & \longrightarrow & A(U_J) & \longrightarrow & 0 \\ p_{J \cup \{k\}}^* \downarrow & & p_J^* \downarrow & & = \downarrow & & \\ \oplus_{k \notin J} A(Y'_{J \cup \{k\}}) & \longrightarrow & A(Y'_J) & \longrightarrow & A(U_J) & \longrightarrow & 0. \end{array}$$

one finds, for each $\alpha'_J \in A(Y'_J)$, classes $\alpha_J \in A(Y_J)$ and $\alpha'_{J \cup \{k\}} \in A(Y'_{J \cup \{k\}})$ such that

$$\alpha'_J = p_J^* \alpha_J + \sum_{k \notin J} \phi_{J*}^{J \cup \{k\}} \alpha'_{J \cup \{k\}}.$$

Reiterating this argument one finds a collection of classes $[\alpha_I] \in \oplus_I A(Y_I)$ for all I containing J , such that

$$\alpha'_J = \sum_{I \supseteq J} p_J^* \phi_{J*}^I \alpha_I.$$

The choice of $[\alpha_I]$ is unique up to the equivalence relation $\sim$. Indeed, if $\sum_{I \supseteq J} p_J^* \phi_{J*}^I \beta_I = 0$ then $\sum_{I \supseteq J} \phi_{J*}^I \beta_I = 0$ by Corollary 1.15, or, equivalently, $\beta_J = -\sum_{I \supset J} \phi_{J*}^I \beta_I$. After applying the equivalence relation to each $\phi_{J*}^I \beta_I$, β_J can then be replaced by a sum of classes from $A(Y_I)$ with $I \supset J$, and induction on the index set can then be applied by Lemma 3.13 in [MM1]. When $J = \emptyset$, the collection $[\alpha_I] \in \oplus_I A(Y_I)$ defines the desired inverse of F .

□

2. THE CHERN CLASSES OF A WEIGHTED PROJECTIVE BLOW-UP

Let $i : Y \hookrightarrow X$ be a regular local embedding of stacks, and consider a groupoid presentation $R \rightrightarrows U$ of X such that $Y \times_X U = \bigsqcup_i V_i$ has all the properties in 1.2. The blow-up $\tilde{X} = \text{Bl}_Y X$ of X along Y has groupoid presentation

$$\tilde{X} = [\text{Bl}_{\{V_i \times_Y V_i\}_i} R \rightrightarrows \text{Bl}_{\{V_i\}_i} U],$$

where the blow-ups are taken in the same fixed order on the étale cover as on relations. If $p : X' \rightarrow X$ is the universally closed étale cover constructed in Section 1, there is a Cartesian diagram

$$\begin{array}{ccc} \tilde{X}' & \xrightarrow{f'} & X' \\ \tilde{p} \downarrow & & \downarrow p \\ \tilde{X} & \xrightarrow{f} & X, \end{array}$$

where $\tilde{X}'$ is obtained by blowing-up X' along the regularly embedded Y'_i in the same order as above. As

$$p^* \mathcal{T}_X \cong \mathcal{T}_{X'}, \quad \tilde{p}^* \mathcal{T}_{\tilde{X}} \cong \mathcal{T}_{\tilde{X}'}, \quad p_i^* \mathcal{N}_{Y|X} \cong \mathcal{N}_{Y'_i|X'}, \quad q_i^* \mathcal{N}_{E|\tilde{X}} \cong \mathcal{N}_{E'_i|\tilde{X}'}$$

for $p_i : Y'_i \rightarrow Y$, and the map between exceptional divisors $q_i : E'_i \rightarrow E$, calculating the Chern invariants of $\tilde{X}$ reduces to the calculation for $\tilde{X}'$ due to Corollary 1.15. We can thus reduce the problem for a regular local embedding to that for a succession of regular embeddings. If, moreover, the ideals $\mathcal{I}_i$ of Y'_i have compatible filtrations with weights such that weighted blow-ups can be defined, then the above reasoning applies to weighted blow-ups as well.

2.1. Weighted projective blow-up and locally trivial weighted projective fibration. Let Y be a regularly embedded substack of the stack X . Let $\pi : \tilde{X} \rightarrow X$ be a weighted blow-up of X along Y , namely,

$$\tilde{X} = \text{Proj}(\oplus_{n \geq 0} \mathcal{I}_n)$$

for an increasing filtration $\{\mathcal{I}_n\}_{n \geq 0}$ of the ideal $\mathcal{I}_Y$ of Y in X , such that $\mathcal{I}_0 = \mathcal{O}_X$, $\mathcal{I}_1 = \mathcal{I}_Y$ and $\mathcal{I}_n \mathcal{I}_m \subseteq \mathcal{I}_{m+n}$ for all $m, n \geq 0$. In addition, assume that the filtration $\{\mathcal{I}_n\}_{n \geq 0}$ satisfies the following properties ([MM1], section 3),

- (1) $\mathcal{I}_k \cap \mathcal{I}_Y^2 = \sum_{j=1}^{k-1} \mathcal{I}_j \mathcal{I}_{k-j}$,
- (2) $\mathcal{I}_k / (\mathcal{I}_k \cap \mathcal{I}_Y^2)$ is a subbundle of the conormal bundle $\mathcal{I}_Y / \mathcal{I}_Y^2$.

such that, if X is smooth then the weighted blow-up stack $\tilde{X}$ is also smooth.

Then the reduced structure of the exceptional divisor in $\tilde{X}$ is

$$E = \text{Proj}(\oplus_{n \geq 0} \mathcal{I}_n / \mathcal{I}_{n+1}),$$

and $\mathcal{I}_n := \pi_* \mathcal{I}_E^n$. Thus E is a locally trivial weighted projective fibration. We recall the following (Lemma 3.1 in [MM1]) description of the Chow ring of E .

Lemma 2.1. *a) The normal bundle in $A := \text{Spec}(\oplus_{n \geq 0} \mathcal{I}_n / \mathcal{I}_{n+1})$ of the fixed locus Y under the natural $\mathbb{C}^*$ action on A is*

$$\mathcal{N}_{Y|A} = \oplus_{n \geq 1} \mathcal{N}_n / \mathcal{N}_{n+1},$$

where $\{\mathcal{N}_n\}_n$ is the filtration of the normal bundle $\mathcal{N}_{Y|X}$ dual to the filtration $\{\mathcal{I}_n / (\mathcal{I}_n \cap \mathcal{I}_Y^2)\}_n$ of $\mathcal{I}_Y / \mathcal{I}_Y^2$.

b) There is a ring isomorphism

$$A^*(E; \mathbb{Q}) \cong \frac{A^*(Y; \mathbb{Q})[\tau]}{\langle P_{Y|X}(\tau) \rangle},$$

where $P_{Y|X}(t)$ is the top equivariant Chern class of the bundle $\mathcal{N}_{Y|A}$. In particular, the free term of $P_{Y|X}(t)$ is the top Chern class of $\mathcal{N}_{Y|X}$. τ is the first Chern class of $\mathcal{O}_E(1)$.

Conversely, given a locally trivial weighted projective fibration $p : P \rightarrow Y$, with a sheaf $\mathcal{L}$ on P such that $P \cong \text{Proj}(\oplus_n \pi_* \mathcal{L}^n)$, then P can be understood as the exceptional divisor of the following weighted blow-up:

$$\tilde{X} := \text{Spec}(\oplus_n \mathcal{L}^n) \rightarrow X := \text{Spec}(\oplus_n \pi_* \mathcal{L}^n).$$

Lemma 2.1 sets up the context for calculating the Chern classes of the locally trivial weighted projective fibration $p : E \rightarrow Y$ by deforming E to a weighted projective bundle on Y and applying the Euler's sequence from Appendix. Indeed, with the notations above, consider the standard deformation $D = \text{Bl}_{Y \times \{\infty\}}(A \times \mathbb{P}^1) \setminus \tilde{A}$ of A to the total space $N_{Y|A}$ of the normal bundle $\mathcal{N}_{Y|A}$, where $\tilde{A} = \text{Bl}_Y A$ is one of the components of the fibre over ∞ of $\text{Bl}_{Y \times \{\infty\}}(A \times \mathbb{P}^1) \rightarrow \mathbb{P}^1$. Thus the fibre over ∞ of $D \rightarrow \mathbb{P}^1$ is $N_{Y|A}$, and the action of $\mathbb{C}^*$ over $A \times \mathbb{P}^1$, with fixed locus $Y \times \mathbb{P}^1$, induces a natural $\mathbb{C}^*$ -action on D . Moreover, due to Lemma 2.1, a), the quotient of $N_{Y|A}$ by $\mathbb{C}^*$ is a weighted projective bundle, and push-forward by the composition

$$N_{Y|A} / \mathbb{C}^* \hookrightarrow D / \mathbb{C}^* \rightarrow E \times \mathbb{P}^1 \rightarrow E$$

induces an isomorphism between the Chow rings of E and of the weighted projective bundle $N_{Y|A} / \mathbb{C}^*$. We obtain the following

Proposition 2.2. *Let $p : P \rightarrow Y$ be a projective fibration as above, and $Q_n := \mathcal{N}_{w_n} / \mathcal{N}_{w_n+1}$, for all indices w_n such that $\mathcal{N}_{w_n} \neq \mathcal{N}_{w_n+1}$, on which $\mathbb{C}^*$ acts with weight w_n . Then the total Chern class*

$$c(P) = p^* c(Y) \prod_n c(Q_n \otimes \mathcal{L}^{\otimes w_n}),$$

with $\mathcal{L} := \mathcal{O}_P(1)$ and $c(Q_n \otimes \mathcal{L}^{\otimes w_n}) := \prod_i (1 + a_i + w_n c_1(\mathcal{L}))$ where a_i are the Chern roots of Q_n .

Proof. Let $i_\infty : N_{Y|A} / \mathbb{C}^* \hookrightarrow D / \mathbb{C}^*$ and $i_0 : P \rightarrow D / \mathbb{C}^*$ be embeddings of fibres in the flat family $D / \mathbb{C}^* \rightarrow \mathbb{P}^1$ and let $q : D / \mathbb{C}^* \rightarrow P$ be the natural projection obtained after taking quotients of $D \rightarrow A$, such that $q \circ i_0 = \text{id}_P$. Then by projection formula and the rational equivalence of fibres,

$$i_{0*} c(\mathcal{T}_P) = i_{0*} i_0^* \mathcal{T}_{D/\mathbb{C}^*} = i_{\infty*} i_\infty^* \mathcal{T}_{D/\mathbb{C}^*} = i_{\infty*} c(N_{Y|A} / \mathbb{C}^*),$$

and thus after composing with q_* ,

$$c(\mathcal{T}_P) = q_* i_{\infty*} c(N_{Y|A} / \mathbb{C}^*),$$

which by Appendix and Lemma 2.1 is of the form described in this Proposition. $\square$

2.2. Model for a weighted blow-up. We start our Chern class calculations with the most approachable type of weighted blow-up: when the blow-up locus is the fixed locus of a $\mathbb{C}^*$ -action on the entire space. In this case, equivariant cohomology techniques permit the recovery of Chern classes of the blow-up from their pull-backs to the exceptional divisor, which in turn are easily computable.

Let Y be a stack, A a $\mathbb{C}^*$ -equivariant affine bundle on Y such that the $\mathbb{C}^*$ -action on A induces a decomposition of the normal bundle of the fixed locus Y

$$\mathcal{N}_{Y|A} = \oplus_h \mathcal{Q}_n$$

with weights $\{w_n\}_n$ and $\text{rk } \mathcal{Q}_n = k_n$. Let $\mathcal{O}_Y$ denote the trivial line bundle on Y . Consider the torus $T := \mathbb{C}^* \times \mathbb{C}^*$ action on $A \times \mathcal{O}_Y$ coming from the individual action of the first $\mathbb{C}^*$ on A and the second on $\mathcal{O}_Y$. In this subsection we will denote by $X := \mathbb{P}^w(A \oplus \mathcal{O}_Y)$, the locally trivial weighted projective fibration obtained as a quotient of $A \times \mathcal{O}_Y$ by $\mathbb{C}^*$ embedded diagonally in T , by $\tilde{Y} := \mathbb{P}^w(A)$, and by $\tilde{X} := \mathbb{P}(\mathcal{O}_{\tilde{Y}}(-1) \oplus \mathcal{O}_{\tilde{Y}})$.

We obtain a blow-up diagram

$$\begin{array}{ccc} \tilde{Y} & \xrightarrow{j} & \tilde{X} \\ g \downarrow & & \downarrow f \\ Y & \xrightarrow{i} & X, \end{array}$$

where the embedding $Y = \mathbb{P}(\mathcal{O}_Y) \hookrightarrow \mathbb{P}^w(A \oplus \mathcal{O}_Y)$, similarly $j : \tilde{Y} \hookrightarrow \tilde{X}$ is the embedding $\tilde{Y} = \mathbb{P}(\mathcal{O}_{\tilde{Y}}) \hookrightarrow \mathbb{P}(\mathcal{O}_{\tilde{Y}}(-1) \oplus \mathcal{O}_{\tilde{Y}})$, the exceptional divisor in the weighted projective blow-down $f : \tilde{X} \rightarrow X$.

Proposition 2.3. *Keeping notations from above, assume that for each n , the total Chern class $c(\mathcal{Q}_n) = c_{k_n}(\mathcal{Q}_n) + \dots + c_1(\mathcal{Q}_n) + 1$ can be written as the pullback of a class $p(\mathcal{Q}_n) = p_{k_n}(\mathcal{Q}_n) + \dots + p_1(\mathcal{Q}_n) + 1 \in H^*(X; \mathbb{Q})$.*

$$c(\tilde{X}) = f^*c(X) \frac{(E+1) \prod_{n=1}^l p(\mathcal{Q}_n(-w_n E))}{\prod_{n=1}^l p(\mathcal{Q}_n)},$$

where E is the class of the exceptional divisor and $p(\mathcal{Q}_n(w_n s)) := \prod_{i=1}^{k_n} (a_i + w_n s + 1)$ where the pullbacks of a_i on Y are the Chern roots of $\mathcal{Q}_n$.

Proof. The morphism f is equivariant with respect to the natural $\mathbb{C}^* := T/\mathbb{C}^*$ -actions on $\tilde{X}$ and X with the weights specified above, such that $\tilde{Y}$ and the section at infinity are the fixed loci for the $\mathbb{C}^*$ -action on $\tilde{X}$, and Y and the section at infinity are the fixed loci for the $\mathbb{C}^*$ -action on X . Thus the diagram above yields another weighted blow-up diagram

$$\begin{array}{ccc} \tilde{Y}_{\mathbb{C}^*} & \xrightarrow{j_{\mathbb{C}^*}} & \tilde{X}_{\mathbb{C}^*} \\ g_{\mathbb{C}^*} \downarrow & & \downarrow f_{\mathbb{C}^*} \\ Y_{\mathbb{C}^*} & \xrightarrow{i_{\mathbb{C}^*}} & X_{\mathbb{C}^*}, \end{array}$$

where $Z_{\mathbb{C}^*} := Z \times_{\mathbb{C}^*} E\mathbb{C}^*$ for each stack Z , and $B\mathbb{C}^*$ is the classifying space of $\mathbb{C}^*$, with universal family $E\mathbb{C}^*$. As noticed above $Y_{\mathbb{C}^*} \cong Y$ and $\tilde{Y}_{\mathbb{C}^*} \cong \tilde{Y}$.

In the following, for easiness of notation, we will drop the subscript $\mathbb{C}^*$ for maps, and only employ it to denote equivariant classes.

In the equivariant cohomology ring $H_{\mathbb{C}^*}^*(\tilde{Y})$

$$\begin{aligned} e^{\mathbb{C}^*}(j^*(\mathcal{T}_{\tilde{X}})) &= e^{\mathbb{C}^*}(\tilde{Y})e^{\mathbb{C}^*}(\mathcal{N}_{\tilde{Y}|\tilde{X}}), \quad e^{\mathbb{C}^*}(i^*(\mathcal{T}_X)) = e^{\mathbb{C}^*}(Y)e^{\mathbb{C}^*}(\mathcal{N}_{Y|X}), \\ \Rightarrow \frac{j^*e^{\mathbb{C}^*}(\tilde{X})}{j^*f^*e^{\mathbb{C}^*}(X)} &= \frac{e^{\mathbb{C}^*}(\mathcal{N}_{\tilde{Y}|\tilde{X}})e^{\mathbb{C}^*}(\tilde{Y})}{e^{\mathbb{C}^*}(\mathcal{N}_{Y|X})e^{\mathbb{C}^*}(Y)} = \frac{(-\xi + t + 1)\prod_{n=1}^l c(\mathcal{Q}_n(w_n\xi))}{\prod_{n=1}^l c(\mathcal{Q}_n(w_nt))}, \end{aligned}$$

where ξ is the first Chern class of $\mathcal{O}_{\tilde{Y}}(1)$.

Let $\alpha := e^{\mathbb{C}^*}(\tilde{X})/f^*e^{\mathbb{C}^*}(X) - 1$ and $P(t) := \frac{(-\xi+t+1)\prod_{n=1}^l p(\mathcal{Q}_n(w_n\xi))}{\prod_{n=1}^l p(\mathcal{Q}_n(w_nt))} - 1$. We note that $P(\xi) = 0$. Thus $\beta := P(t)/(t - \xi)$ is well defined and by the self intersection formula on the exceptional divisor,

$$(2.1) \quad j^*\alpha = j^*j_*\beta.$$

By the Atiyah-Bott localization theorem,

$$(2.2) \quad \alpha = j_* \frac{j^*\alpha}{e^{\mathbb{C}^*}(\mathcal{N}_{\tilde{Y}|\tilde{X}})} + l_* \frac{l^*\alpha}{e^{\mathbb{C}^*}(\mathcal{N}_{E_\infty|\tilde{X}})},$$

where $j : \tilde{Y} \hookrightarrow \tilde{X}$ and $l : E_\infty \hookrightarrow \tilde{X}$ are the two fixed point loci of $\tilde{X}$ under the $\mathbb{C}^*$ -action. On the one hand, by relation 2.1

$$j_* \frac{j^*\alpha}{e^{\mathbb{C}^*}(\mathcal{N}_{\tilde{Y}|\tilde{X}})} = j_* \frac{j^*j_*\beta}{e^{\mathbb{C}^*}(\mathcal{N}_{\tilde{Y}|\tilde{X}})} = j_*\beta$$

and on the other hand, as the section at infinity $E_\infty = \mathbb{P}^w(N)$ and $\tilde{Y}$ are disjoint, $l^*\mathcal{T}_{\tilde{X}} = l^*f^*\mathcal{T}_X$ and thus the second term on the right hand side of equation 2.2 is zero. Thus in fact $\alpha = j_*\beta$ which implies

$$e^{\mathbb{C}^*}(\tilde{X}) = \frac{(E + t + 1)\prod_{n=1}^l p(\mathcal{Q}_n(-w_nE))}{\prod_{n=1}^l p(\mathcal{Q}_n(w_nt))}.$$

The classical limit $t \rightarrow 0$ yields the desired relation. □

2.3. Deformation to the weighed normal cone. Returning to the general case, let Y be a regularly embedded substack of the stack X . Let $\pi : \tilde{X} \rightarrow X$ be the weighted blow-up of X along Y for an increasing filtration $\{\mathcal{I}_n\}_{n \geq 0}$ of the ideal $\mathcal{I}_Y$, satisfying properties (1) and (2) in 2.1. The ideal sheaf $\mathcal{J}$ of $Y \times \{\infty\}$ in $X \times \mathbb{P}^1$ admits a filtration formed by the sheaves $\mathcal{J}_n := \sum_{k=0}^n \mathcal{I}_k \mathcal{K}^{n-k}$ where $\mathcal{K}$ is the ideal of ∞ in $\mathbb{P}^1$ pulled back to $X \times \mathbb{P}^1$. Let $M = \text{Proj}(\oplus_n \mathcal{J}_n)$ be the weighted projective blow-up of $X \times \mathbb{P}^1$ along $Y \times \{\infty\}$ with the filtration $\{\mathcal{J}_n\}_n$.

The usual properties of the deformation to the normal cone carry out for this construction with the suitable changes in weights:

(1) There is a natural closed regular embedding $J : Y \times \mathbb{P}^1 \hookrightarrow M$.

(2) The composition $\rho = p_2 \circ P$ of the blow-up map $P : M \rightarrow X \times \mathbb{P}^1$ with the projection $p_2 : X \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is a flat morphism of stacks, and the following diagram

commutes

$$\begin{array}{ccc} Y \times \mathbb{P}^1 & \xrightarrow{J} & M \\ \text{\scriptsize pr_2} \downarrow & \swarrow \rho & \\ \mathbb{P}^1 & & \end{array}$$

(3) Over $\mathbb{P}^1 \setminus \{\infty\}$, the preimage $\rho^{-1}(\mathbb{A}^1) = X \times \mathbb{A}^1$ and J is the trivial embedding.

(4) As a Cartier divisor,

$$M_\infty := \rho^{-1}(\infty) = P + \tilde{X},$$

where $P = \text{Proj}(\oplus_{n \geq 0} \mathcal{I}_n / \mathcal{I}_{n+1})$ is a locally trivial weighted projective fibration and $\tilde{X} = \text{Proj}(\oplus_{n \geq 0} \mathcal{I}_n)$ is the weighted blow-up of X along the locus Y . Both P and $\tilde{X}$ are Cartier divisors of M , intersecting in $E := \text{Proj}(\oplus_{n \geq 0} \mathcal{I}_n / \mathcal{I}_{n+1})$, which is embedded as the section at infinity in P and as the exceptional divisor in $\tilde{X}$. On the other hand, $Y_\infty = Y \times \{\infty\}$ embeds in M_∞ as the zero section in P and is thus disjoint from $\tilde{X} \hookrightarrow M_\infty$.

The proof is analogous to [F], (5.1).

Theorem 2.4. *Consider the weighted blow-up $f : \tilde{X} \rightarrow X$ of a stack X along a regularly embedded substack Y with an increasing filtration $\{\mathcal{I}_n\}_{n \geq 0}$ of the ideal $\mathcal{I}_Y$, satisfying conditions (1)-(2) in 2.1. Let $\{\mathcal{N}_n\}_n$ denote the filtration of the normal bundle $\mathcal{N}_{Y|X}$ dual to the filtration $\{\mathcal{I}_n / (\mathcal{I}_n \cap \mathcal{I}_Y^2)\}_n$ of $\mathcal{I}_Y / \mathcal{I}_Y^2$. Let $\mathcal{Q}_n = \mathcal{N}_n / \mathcal{N}_{n+1}$.*

Assume that for each n , the total Chern class $c(\mathcal{Q}_n) = c_{k_n}(\mathcal{Q}_n) + \dots + c_1(\mathcal{Q}_n) + 1$ can be written as the pullback of a class $p(\mathcal{Q}_n) = p_{k_n}(\mathcal{Q}_n) + \dots + p_1(\mathcal{Q}_n) + 1 \in H^(X; \mathbb{Q})$. Then*

$$c(\tilde{X}) = f^*c(X) \frac{(E+1) \prod_{n=1}^l p(\mathcal{Q}_n(-w_n E))}{\prod_{n=1}^l p(\mathcal{Q}_n)},$$

where E is the class of the exceptional divisor in $\tilde{X}$ and $p(\mathcal{Q}_n(w_n s)) := \prod_{i=1}^{k_n} (a_i + w_n s + 1)$ where the pullbacks of a_i on Y are the Chern roots of $\mathcal{Q}_n$.

Proof. Let $\tilde{M}$ be the weighted blow-up of M along $Y \times \mathbb{P}^1$ with the filtration $r^* \mathcal{I}_n$, where $r = p_1 \circ P$ is the composition $M \rightarrow X \times \mathbb{P}^1 \rightarrow X$. Looking at the fibres over 0 and $\infty \in \mathbb{P}^1$

$$\begin{array}{ccccc} \tilde{X} & \xrightarrow{\tilde{j}_0} & \tilde{M} & \xleftarrow{\tilde{j}_\infty} & \tilde{P} + \tilde{X} \\ f_0 \downarrow & & \downarrow F & & \downarrow f_\infty \\ X & \xrightarrow{j_0} & M & \xleftarrow{j_\infty} & P + \tilde{X} \\ \downarrow & & \downarrow \rho & & \downarrow \\ 0 & \longrightarrow & \mathbb{P}^1 & \longleftarrow & \infty \end{array}$$

The embeddings $\tilde{j}_\infty$, respectively j_∞ split into $\tilde{k} : \tilde{P} \rightarrow \tilde{M}$, $\tilde{l} : \tilde{X} \rightarrow \tilde{M}$, respectively $k : P \rightarrow M$, $l : \tilde{X} \rightarrow M$, where $F \circ \tilde{l}$ and l can be naturally identified. Pullbacks of the quotient sheaf $\mathcal{G}_M := \mathcal{T}_{\tilde{M}} / F^* \mathcal{T}_M$ yield $\tilde{j}_0^* \mathcal{G}_M = \mathcal{G}_0 := \mathcal{T}_{\tilde{X}} / f_0^* \mathcal{T}_X$ and $\tilde{j}_\infty^* \mathcal{G}_M = \mathcal{G}_\infty$ which

on $\tilde{P}$ is the quotient $\mathcal{T}_{\widetilde{\mathbb{P}^w(C \oplus 1)}}/f^*\mathcal{T}_P$, and on $\tilde{X}$ is the zero sheaf. We note that the maps of locally free sheaves on stacks

$$\mathcal{T}_{\tilde{M}} \rightarrow F^*\mathcal{T}_M, \quad \mathcal{T}_{\tilde{X}} \rightarrow f_0^*\mathcal{T}_X \quad \text{and} \quad \mathcal{T}_{\tilde{P}} \rightarrow f_\infty^*\mathcal{T}_P$$

are monomorphisms, as $\tilde{M}$ and M are isomorphic outside the exceptional divisor and its image, etc. Moreover,

$$\tilde{j}_{0*}c(\mathcal{G}_0) = \tilde{j}_{0*}\tilde{j}_0^*c(\mathcal{G}_M) = c(\mathcal{G}_M) \cdot [\tilde{M}_0],$$

which by rational equivalence is identified with

$$c(\mathcal{G}_M) \cdot [\tilde{M}_\infty] = \tilde{j}_{\infty*}c(\mathcal{G}_\infty) = k_*c(\mathcal{G}_{\infty|\tilde{P}}) + [\tilde{X}],$$

as $\tilde{l}^*c(\mathcal{G}_M) = c(\tilde{l}^*\mathcal{G}_M) = 1$ on $\tilde{X}$. On the one hand, the formula for $c(\mathcal{G}_{\infty|\tilde{P}})$ is given by ??.

On the other hand, $\tilde{M}$ is isomorphic to the blow-up of $\tilde{X} \times \mathbb{P}^1$ along $E \times \{\infty\}$, where E is the exceptional divisor of $\tilde{X}$. As such, there is a map $\tilde{P} : \tilde{M} \rightarrow \tilde{X} \times \mathbb{P}^1$ whose composition q with the projection $\tilde{X} \times \mathbb{P}^1 \rightarrow \tilde{X}$, satisfies $q \circ \tilde{j}_0 = \text{id}_{\tilde{X}}$. From this and the model case in Proposition 2.3 we recover the general formula in the theorem. □

3. THE MODULI SPACE $\overline{M}'_{0,m}(\mathbb{P}^n, d)$

3.1. Let m, n, d be nonnegative integers. In this section we apply the general theory for weighted blow-ups along regular local embeddings to calculate the total Chern class of the moduli space of stable maps $\overline{M}_{0,m}(\mathbb{P}^n, d)$. We first set up the context by briefly recalling the blow-up constructions of $\overline{M}_{0,m}(\mathbb{P}^n, d)$ from [MM1] (for $m = 1$), [MM2] (for $m > 1$), and [10] (for $m = 0$). They pertain to a family of smooth Deligne-Mumford stacks $\overline{M}_{0,\mathcal{A}}(\mathbb{P}^n, d, a)$, and of weighted blow-ups

$$\overline{M}_{0,\mathcal{A}}(\mathbb{P}^n, d, a) \rightarrow \overline{M}_{0,\mathcal{A}'}(\mathbb{P}^n, d, a'),$$

where $a, a' \in \mathbb{Q}$, $\mathcal{A} = (a_1, \dots, a_m)$, $\mathcal{A}' = (a'_1, \dots, a'_m) \in \mathbb{Q}^m$ such that

$$\sum_{i=1}^m a_i + da > 2, \quad \sum_{i=1}^m a'_i + da' > 2, \quad a \geq a' > 0, \quad 1 \geq a_j \geq a'_j \geq 0$$

for all $j = 1, \dots, m$. Here $\overline{M}_{0,\mathcal{A}}(\mathbb{P}^n, d, a)$ is the stack of $(\mathcal{A}, a)$ -weighted stable maps as defined in [MM2], parametrizing families of rational curves $\pi : C \rightarrow S$, whose fibers are either smooth or with nodes as singularities, with m marked sections not intersecting the nodes of the fibers, a line bundle $\mathcal{L}$ on C of degree d on each fiber C_s , and a morphism $e : \mathcal{O}_C^{n+1} \rightarrow \mathcal{L}$ (specified up to isomorphisms of the target) satisfying a series of stability conditions:

- (1) $\omega_{C|S}(\sum_{i=1}^m a_i p_i) \otimes \mathcal{L}^a$ is relatively ample over S ,
- (2) $\mathcal{G} := \text{coker } e$, restricted over each fiber C_s , is a skyscraper sheaf supported only on smooth points of C_s , and
- (3) for any $s \in S$ and $p \in C_s$ and for any $I \subseteq \{1, \dots, m\}$ (possibly empty) such that $p = p_i$ for all $i \in I$ the following holds

$$\sum_{i \in I} a_i + a \dim \mathcal{G}_p \leq 1.$$

Let C be a curve. A tail of C is a closed connected subcurve C' of C with the property that $C \setminus C'$ is connected.

Definition 3.1. $I \subset \mathcal{P}(D) \setminus \{\emptyset, D\}$ is a nested set if, for any two $h, h' \in I$, the intersection $h \cap h'$ is either h , h' or $\emptyset$.

Definition 3.2. Fix a positive number $k < d$ and a nested set $I \subset \mathcal{P} \setminus \{\emptyset, D\}$ such that $h \cap h' = \emptyset$ for any distinct $h, h' \in I_{\leq d-k}$.

$\overline{M}_I^k$ is defined in [MM1], (Proposition 2.3) as the stack of I -type, k -stable, degree d maps from a rational curve into $\mathbb{P}^n$, i.e.

$$(C, p_1, \{p_h\}_{h \in I_{\leq d-k}}, \{C_h\}_{h \in I_{> d-k}}, \mathcal{L}, e)$$

of a k -stable, degree d pointed map $(C, p_1, \mathcal{L}, e)$, together with marked points $\{p_h\}_{h \in I_{\leq d-k}}$ and connected subcurves $\{C_h\}_{h \in I_{> d-k}}$ satisfying the following properties:

- (1) $\forall h \in I_{> d-k}$, $p_1 \notin C_h \subset C$ and $\deg \mathcal{L}|_{C_h} = |h|$;
- (2) $\forall h \in I_{\leq d-k}$, $\dim \operatorname{coker} e_{p_h} = |h|$;
- (3) compatibility of incidence relations:
 - $\forall h \in I_{\leq d-k}$, $\forall h' \in I_{> d-k}$, $h \subset h'$ iff $p_h \in C_{h'}$;
 - $\forall h, h' \in I_{> d-k}$, if $h' \subset h$, then $C_{h'} \subset C_h$, if $h \subset h'$, then $C_h \subset C_{h'}$, otherwise $C_h \cap C_{h'} = \emptyset$;

A curve C which admits a set of points and components with the above properties is said to be of I -splitting type.

We note that while for the generic curve parametrized by $\overline{M}_I^k$, all tails and base locus points are marked by elements of I , other curves, represented by points in the boundary of $\overline{M}_I^k$, may have unmarked tails and base points. It is due to such points that the maps $\overline{M}_J^k \rightarrow \overline{M}_I^k$ are in general only local embedding. This suggests that marking all components and base points of curves and their maps will result in a moduli problem for the stack $\overline{M}_{0,m}(\mathbb{P}^m, d)'$ as in section 1.

Definition 3.3. Consider the moduli functor from schemes to sets, associating to any scheme S the set $(\mathcal{A}, a)$ -stable, degree d pointed maps $(C \rightarrow S, \{p_i\}_{i \in \{1, \dots, m\}}, \mathcal{L}, e)$, together with a collection $\{P_s\}_{s \in S}$ of partitions

$$D = \sqcup_{\alpha \in N_s} B_\alpha$$

of the set $D = \{1, \dots, d\}$, one for each $s \in S$, such that for every $s \in S$, the set

$$N_s \cong \{ \text{irreducible components of the curve } C_s \} \sqcup \{x \in C_s \mid \operatorname{coker} e_x \neq \emptyset\},$$

and the partition is compatible with the structure of the map given by $(\mathcal{L}, e)$ in the following sense

- (1) If $\alpha \in N_s$ corresponds to the component $C \subset C_s$ then $|B_\alpha| = \deg \mathcal{L}|_C$;
- (2) If $\alpha \in N_s$ corresponds to the point $x \in C_s$ then $|B_\alpha| = \dim \operatorname{coker} e_x$.

A set of data as above will be called a semi-rigid $(\mathcal{A}, a)$ -stable map over S . An isomorphism of semi-rigid $(\mathcal{A}, a)$ -stable maps is an isomorphism of $(\mathcal{A}, a)$ -stable maps which also preserves the partitions of D .

Let $\overline{M}_{0,\mathcal{A}}(\mathbb{P}^m, d, a)'$ be the stack associated to $\overline{M}_{0,\mathcal{A}}(\mathbb{P}^m, d, a)'$ and all the local embeddings $\overline{M}_I^a \rightarrow \overline{M}_{0,\mathcal{A}}(\mathbb{P}^m, d, a)$, like in Theorem 1.9.

Theorem 3.4. *The moduli problem of semi-rigid $(\mathcal{A}, a)$ -stable maps is finely represented by the Deligne-Mumford stack $\overline{M}'_{0,\mathcal{A}}(\mathbb{P}^n, d, a)$.*

Proof. For any semi-rigid $(\mathcal{A}, a)$ -stable map $(C \rightarrow S, \{p_i\}_{i \in \{1, \dots, m\}}, \mathcal{L}, e, \{P_s\}_{s \in S})$ a map $S \rightarrow \overline{M}_{0,\mathcal{A}}(\mathbb{P}^m, d, a)$ is obtained by forgetting the D -partitions. The partition P_s on each fiber C_s over $s \in S$ completely determines the splitting types of C_s and thus the fiber product $S_I := S \times_{\overline{M}_{0,\mathcal{A}}(\mathbb{P}^m, d, a)} \overline{M}_I^a$ is embedded in S , for each I as above. By the universal property 1.5 of $\overline{M}'_{0,\mathcal{A}}(\mathbb{P}^n, d, a)$, there is a map

$$S \rightarrow \overline{M}'_{0,\mathcal{A}}(\mathbb{P}^n, d, a)$$

commuting with the map to $\overline{M}_{0,\mathcal{A}}(\mathbb{P}^n, d, a)$. As such, $(C \rightarrow S, \{p_i\}_{i \in \{1, \dots, m\}}, \mathcal{L}, e)$ is the pullback of the $(\mathcal{A}, a)$ -stable map structure of the universal family on $\overline{M}'_{0,\mathcal{A}}(\mathbb{P}^n, d, a)$. As for every $s \in S$, the partition P_s on C_s is completely determined by the set $\{I; s \in S_I\}$, it follows that the partition P_s is also inherited by pullback from the universal family on $\overline{M}'_{0,\mathcal{A}}(\mathbb{P}^n, d, a)$. □

Thus for $m \geq 1$, the moduli space of stable maps with marked points $\overline{M}_{0,m}(\mathbb{P}^n, d)$ corresponds to weights $a > 1$ and $\mathcal{A} = (1, 1, \dots, 1)$, and a sequence of birational contractions of $\overline{M}_{0,m}(\mathbb{P}^n, d)$ is given by $\overline{M}_{0,\mathcal{A}_k}(\mathbb{P}^n, d, a_k)$ with $a_k = \frac{1}{k+\epsilon}$ for $0 < \epsilon < 1$ and $\mathcal{A}_k = (1, a_k, \dots, a_k)$ for $d > k > 0$. When $m = 1$, the last member in the sequence of contractions, $\overline{M}_{0,1}(\mathbb{P}^n, d, a_{d-1})$ is a weighted projective fibration $\mathbb{P}(A)$ over $\mathbb{P}^n$ described in [MM1], Lemma 3.3.

The normal bundle of the zero section $\mathcal{N}_{\mathbb{P}^n|A} = \oplus_k \mathcal{N}_k / \mathcal{N}_{k+1}$, where

$$\mathcal{N}_k = (\oplus_{i=1}^{(n+1)(d-k)} \mathcal{O}_{\mathbb{P}^n}(1)) / \mathcal{O}_{\mathbb{P}^n}.$$

$\mathbb{C}^*$ acts on the bundle $\mathcal{N} = \oplus_k \mathcal{N}_k / \mathcal{N}_{k+1}$ with weights $(1, \dots, d)$ and thus by ..., the total Chern class

$$c(\overline{M}_{0,1}(\mathbb{P}^n, d, a_{d-1})) = (1 + H)^{n+1} (1 + \psi)^{-1} \prod_{k=1}^d (1 + H + k\psi)^{n+1}.$$

More generally for any I , denote $l_I := |\cup_{h \in I} h|$ and s_I be the number of maximal elements of I , i.e. elements $h \in I$ such that there is no $h' \in I$ with $h \subset h'$.

The normal bundle

$$\mathcal{N}_{\overline{M}_I^{a_{d-1}}|\overline{M}^{a_{d-1}}} = \mathcal{N}_{\mathbb{P}^1 \times \mathbb{P}^n | (\mathbb{P}^1)^{s_I} \times \mathbb{P}_{d-l_I}^n} \Big|_{\mathbb{P}^n} = (\mathcal{O}_{\mathbb{P}^n}^{s_I} \oplus \oplus_{i=1}^{(n+1)(d-l_I)} \mathcal{O}_{\mathbb{P}^n}(1)) / \mathcal{O}_{\mathbb{P}^n}$$

admits a natural filtration $\{\mathcal{N}_k\}_k$ described in Lemma 3.2. Here

$$\mathcal{N}_k = (\mathcal{O}_{\mathbb{P}^n}^{s_I} \oplus \oplus_{i=1}^{(n+1)(d-l_I-k)} \mathcal{O}_{\mathbb{P}^n}(1)) / \mathcal{O}_{\mathbb{P}^n}$$

$\mathbb{C}^*$ acts on the bundle $\mathcal{N} = \oplus_k \mathcal{N}_k / \mathcal{N}_{k+1}$ with weights $(1, \dots, d - l_I)$. Thus according to formula ... the total Chern class of the weighted projective fibration $\overline{M}_I^{a_{d-1}}$ over $\mathbb{P}^n$ is

$$c(\overline{M}_I^{a_{d-1}}) = (1 + H)^{n+1} (1 + \psi)^{s_I-1} \prod_{i=1}^{d-l_I} (1 + H + i\psi)^{n+1}.$$

The normal bundle of the blow-up locus for $\overline{M}_I^{a_{k-1}} \rightarrow \overline{M}_I^{a_k}$ when $k \in \{1, \dots, d-1\}$ and the weights of the appropriate $\mathbb{C}^*$ action on it have been calculated in [MM1], Lemma 3.21. Let $h \in \mathcal{P}(D)$ be such that $|h| = k$ and denote by $I_h \subseteq I$ the set of all $h' \in I$ such that $h' \subset h = \emptyset$. The essential ingredients for the Chern class calculations are the classes

$$\psi_h = \psi - \sum_{h' \supseteq h} D_{h'} \text{ and } \psi_h^0 = \psi - \sum_{h' \supset h} D_{h'},$$

$$H_{I;h} = H + (d - |h \cup (\cup_{h' \in I_h} h')|) \psi - \sum_{h' \supset h} |h' \setminus (h \cup (\cup_{h'' \in I} h''))| D_{h'}.$$

The top equivariant Chern class of the normal bundle $\mathcal{N}_{\overline{M}_{I \cup \{h\}}^{a_k} | \overline{M}_I^{a_k}}$ when evaluated at $t = D_h$ is

$$(1 + \psi_h)^{|I_h|-1} \prod_{j=1}^{|h \setminus \cup_{h' \in I_h} h'|} (1 + H_{I;h} + j\psi_h)^{n+1}$$

and thus formula 2.1 reads in this case

$$c(\overline{M}_I^{a_{k-1}}) = f_I^{a_{k-1}*} c(\overline{M}_I^{a_{k-1}}) \prod_{h; |h|=k} \frac{(1 + D_h)(1 + \psi_h)^{|I_h|-1} \prod_{j=1}^{|h \setminus \cup_{h' \in I_h} h'|} (1 + H_{I;h} + j\psi_h)^{n+1}}{(1 + \psi_h^0)^{|I_h|-1} \prod_{j=1}^{|h \setminus \cup_{h' \in I} h'|} (1 + H_{I;h} + j\psi_h^0)^{n+1}}.$$

Iterating the formula above for all $k \in \{1, \dots, d-1\}$,

Theorem 3.5. *The total Chern class of the normalized stratum $\overline{M}_I^{a_k}$ of $\overline{M}^{a_k}$*

$$c(\overline{M}_I^{a_k}) = (1 + H)^{n+1} (1 + \psi)^{s_I-1} \prod_{i=1}^{d-l_I} (1 + H + i\psi)^{n+1} \prod_{h; |h| > k} \frac{(1 + D_h)(1 + \psi_h)^{|I_h|-1} \prod_{j=1}^{|h \setminus \cup_{h' \in I_h} h'|} (1 + H_{I;h} + j\psi_h)^{n+1}}{(1 + \psi_h^0)^{|I_h|-1} \prod_{j=1}^{|h \setminus \cup_{h' \in I} h'|} (1 + H_{I;h} + j\psi_h^0)^{n+1}}$$

where the product is taken after all h such that $I \cup \{h\}$ is still a nested set. $l_I := |\cup_{h \in I} h|$ and s_I is the number of maximal elements of I .

In particular,

$$c(\overline{M}_{0,1}(\mathbb{P}^n, d)) = (1 + H)^{n+1} \frac{\prod_{i=1}^d (1 + H + i\psi)^{n+1}}{(1 + \psi)} \prod_h \frac{(1 + D_h)(1 + \psi_h^0) \prod_{j=1}^{|h \setminus \cup_{h' \in I} h'|} (1 + H_h + j\psi_h)^{n+1}}{(1 + \psi_h) \prod_{j=1}^{|h \setminus \cup_{h' \in I} h'|} (1 + H_h + j\psi_h^0)^{n+1}},$$

where $H_h = H + (d - |h|)\psi - \sum_{h' \supset h} |h' \setminus h| D_{h'}$ is $H_h^\emptyset$ from above. As described in [MM2], the spaces $\overline{M}_{0,m}(\mathbb{P}^n, d, a_k)$ with $k \in \{1, \dots, d-1\}$ can be thought of as normalized substrata of $\overline{M}_{0,1}(\mathbb{P}^n, d + m - 1, a_k)$. More precisely, after choosing a privileged point 1_M among the m marked points, and adding the rest of marked points $M' = \{2_M, \dots, m_M\}$ to the set of component/base locus $\{1_D, \dots, d_D\}$ to form $D' = \{1_D, \dots, d_D, 2_M, \dots, m_M\}$, one can write

$$\overline{M}_{0,m}(\mathbb{P}^n, d, a_k) \cong \overline{M}_I^{a_k}$$

for $I = \{\{2_M\}, \{3_M\}, \dots, \{d_M\}\}$. Thus

$$c(\overline{M}_{0,m}(\mathbb{P}^n, d)) = (1+H)^{n+1}(1+\psi)^{m-2} \prod_{i=1}^d (1+H+i\psi)^{n+1} \\ \prod_{h \in \mathcal{P}(D') \setminus \{\emptyset, \{2_M\}, \dots, \{m_M\}\}} \frac{(1+D_h)(1+\psi_h)^{|h_M|-1} \prod_{j=1}^{|h \setminus \cup_{h' \in I} h'|} (1+H_{m;h}+j\psi_h)^{n+1}}{(1+\psi_h^0)^{|h_M|-1} \prod_{j=1}^{|h \setminus \cup_{h' \in I} h'|} (1+H_{m;h}+j\psi_h^0)^{n+1}},$$

where $h_M = h \cap \{2_M, \dots, m_M\}$ and

$$H_{m;h} = H + |D \setminus h|\psi - \sum_{h' \supset h} |D \cap h' \setminus h| D_{h'}.$$

In particular, one recovers the formulae for the first Chern classes as calculated in [P] and in a more general setup in [DJS]

Let $f : \overline{M}_{0,1}(\mathbb{P}^n, d) \rightarrow \overline{M}_{0,0}(\mathbb{P}^n, d)$ be the forgetful morphism. In [10] this has been split into a sequence of simple blow-ups and a relatively simpler fibration $\overline{U}^{[(d-1)/2]} \rightarrow \overline{M}_{0,0}(\mathbb{P}^n, d)$.

Choose $I := \{h \subset \{1, \dots, d\}, |h| > d/2\}$ if d is odd, and let I additionally contain half of the sets h with $|h| = d/2$ if d is even, under the condition that no two sets $h, \bar{h}$ are simultaneously in I . The class

$$\psi'_I := \psi - \sum_{h \in I} D_h \in B^*(\overline{M}_{0,1}(\mathbb{P}^n, d))$$

can be considered as the $\mathcal{O}(1)$ -line bundle for the fibration above. When d is odd, ψ'_I is the pullback of the relative cotangent class for the morphism $\overline{U}^{[(d-1)/2]} \rightarrow \overline{M}_{0,0}(\mathbb{P}^n, d)$. The classes D_h with $h \in I$ are pullbacks of the classes of the exceptional divisors on $\overline{M}_{0,1}(\mathbb{P}^n, d)$. A formula comparing the total Chern classes,

$$f^*(c(\overline{M}_{0,0}(\mathbb{P}^n, d))) = c(\overline{M}_{0,1}(\mathbb{P}^n, d)) \frac{\prod_{h \in I} (1+D_h) P_h(1-D_h)}{P(1+\psi_I) \prod_{h \in I} P_h(1)}$$

where $P(\psi_I)$ and $P_h(-D_h)$ are the quadratic expressions in [MM2], Theorem 3.3., (1)–(2). Proposition 2.1. in [10] shows how to recover the class $c(\overline{M}_{0,0}(\mathbb{P}^n, d))$ from its pullback.

APPENDIX. EULER'S SEQUENCE FOR A WEIGHTED PROJECTIVE BUNDLE

Let $g : P \rightarrow Y$ be a weighted projective bundle and a smooth morphism of stacks. With the notations from above, consider the splitting $\mathcal{N}_{Y|A} = \oplus_i L_i$, such that for all n with $\mathcal{N}_n \neq \mathcal{N}_{n+1}$ there is a unique index i with $L_i = \mathcal{N}_n / \mathcal{N}_{n+1}$. Denote $w_i = n$ the weight of the naturally induced $\mathbb{C}^*$ -action on L_i .

Let $\mathcal{T}_{P|Y} = \ker(\mathcal{T}_P \rightarrow g^* \mathcal{T}_Y)$ denote the relative tangent bundle of $g : P \rightarrow Y$.

Lemma 3.6. *There is an exact sequence of vector bundles on P*

$$(3.1) \quad 0 \longrightarrow \mathcal{O}_P \xrightarrow{\sigma} \oplus_i g^* L_i \otimes \mathcal{O}_P(w_i) \xrightarrow{e} \mathcal{T}_{P|Y} \longrightarrow 0.$$

Proof. The weighted projective bundle P is locally trivial over Y , i.e. $P|_U \cong \mathbb{P}[w_0 : \dots : w_n] \times U$ for each open set U in some open cover of Y . We first discuss briefly the case

of the weighted projective space $\mathcal{P}[w_0 : \dots : w_n] = [\text{Spec } \mathbb{C}[x_0, \dots, x_n] \setminus \{0\} / \mathbb{C}^*]$, where $\mathbb{C}^*$ acts on x_i with weight w_i .

An étale presentation of the Deligne-Mumford stack is $\mathcal{P}[w_0 : \dots : w_n]$ is given by

$$\bigsqcup_{i,j \in \{0, \dots, n\}} V_{ij} \rightrightarrows \bigsqcup_{i \in \{0, \dots, n\}} V_i$$

where $V_i = \mathbb{C}^n$, $V_{ij} = V_{ji} = \mathbb{C}^{n-1} \times \mathbb{C}^*$ with coordinates $\{v_i^k\}_{k \in \{0, \dots, n\} \setminus \{i\}}$ such that $v_i^j \neq 0$, the étale maps $\phi_{ij}^i : V_{ij} \rightarrow V_i$ given by

$$\phi_{ij}^i(\{v_i^k\}_k) = (v_i^0, \dots, (v_i^j)^{w_j}, \dots, v_i^n),$$

with $(v_i^j)^{w_j}$ in the j -th position, and the change of coordinates

$$v_j^i = \frac{1}{v_i^j}, \quad v_j^k = \frac{v_i^k}{(v_i^j)^{w_k}} \text{ for } k \neq i, j$$

on V_{ij} .

Let $\{u_i^k\}_{k \in \{0, \dots, n\} \setminus \{i\}}$ denote natural coordinates on V_i , such that the map to the coarse moduli space $\mathcal{P}[w_0 : \dots : w_n] \rightarrow \mathbb{P}[w_0 : \dots : w_n]$ sends $(u_i^k)_{k \in \{0, \dots, n\} \setminus \{i\}} \in V_i$ to the point $[u_i^0 : \dots : 1 : \dots : u_i^n]$ with 1 in the i -th position.

The line bundle $\mathcal{O}_{\mathcal{P}[w_0 : \dots : w_n]}(1)$ is determined by the trivialisations $L_i \cong V_i \times \mathbb{C}$, with gluing maps

$$\phi_{ij}^{i*} L_i \rightarrow \phi_{ij}^{j*} L_j$$

given by $\phi_{ij}^{j*} s_j = v_i^j \phi_{ij}^{i*} s_i$, where s_i and s_j are the unitary sections on L_i and L_j . Thus $\mathcal{O}_{\mathcal{P}[w_0 : \dots : w_n]}(1)$ is the pullback of the line bundle $\mathcal{O}_{\mathbb{P}[w_0 : \dots : w_n]}(1)$ on the coarse moduli scheme, whose total space is

$$\mathbb{P}[w_0 : \dots : w_n : d] \setminus \{[0 : \dots : 0 : 1]\} \rightarrow \mathbb{P}[w_0 : \dots : w_n].$$

Here $d = \gcd(w_0, \dots, w_n) = 1$ and the map above is the projection on the first $(n+1)$ coordinates. Each weighted projective coordinate x_i gives rise to a global section of $\mathcal{O}_{\mathbb{P}[w_0 : \dots : w_n]}(w_i)$, whose pullback to $\mathcal{O}_{\mathcal{P}[w_0 : \dots : w_n]}(w_i)$ will be denoted by x_i as well.

Note that the condition $\gcd(w_0, \dots, w_n) = 1$ implies the smoothness of the stack $\mathcal{P}[w_0 : \dots : w_n]$. The tangent bundle to $\mathcal{P}[w_0 : \dots : w_n]$ is determined by the vector bundles $\mathcal{T}_{V_i} \cong V_i \times \mathbb{C}^n$ with isomorphisms

$$\phi_{ij}^{i*} \mathcal{T}_{V_i} \cong \phi_{ij}^{j*} \mathcal{T}_{V_j},$$

identifying

$$\phi_{ij}^{j*} \left(\frac{\partial}{\partial u_j^k} \right) = (v_i^j)^{w_k} \phi_{ij}^{i*} \left(\frac{\partial}{\partial u_i^k} \right),$$

$$\phi_{ij}^{i*} \left(\frac{\partial}{\partial u_j^i} \right) = (v_i^j)^{w_i} \phi_{ij}^{i*} \left(- \sum_{k \neq i} \frac{w_k}{w_i} v_i^k \frac{\partial}{\partial u_i^k} \right).$$

Indeed, $\phi_{ij}^{i*} \mathcal{T}_{V_i} = \langle \left\{ \phi_{ij}^{i*} \left(\frac{\partial}{\partial u_i^k} \right) = \frac{\partial}{\partial v_i^k} \right\}_{k \in \{0, \dots, n\} \setminus \{i\}}, \phi_{ij}^{i*} \left(\frac{\partial}{\partial u_j^i} \right) = \frac{1}{w_j} (v_i^j)^{w_j-1} \frac{\partial}{\partial v_i^j} \rangle$, and the identifications above are naturally derived from the change of coordinates v_i^k into v_j^k .

The short exact sequence

$$(3.2) \quad 0 \longrightarrow \mathcal{O}_{\mathcal{P}[w_0:\dots:w_n]} \xrightarrow{\sigma} \oplus_i \mathcal{O}_{\mathcal{P}[w_0:\dots:w_n]}(w_i) \xrightarrow{e} \mathcal{T}_{\mathcal{P}[w_0:\dots:w_n]} \longrightarrow 0$$

is defined locally by

$$\sigma(1) := (w_0 x_0, \dots, w_n x_n),$$

$$e(s_i^{\otimes a_k}) := \frac{\partial}{\partial u_i^k}, \quad e(s_i^{\otimes a_i}) := - \sum_{k \neq i} \frac{w_k}{w_i} u_i^k \frac{\partial}{\partial u_i^k}$$

and by the presentation of the bundles above this is well defined globally.

Given a weighted projective bundle $g : P = [A/\mathbb{C}^*] \rightarrow Y$ with weights $(w_0, \dots, w_n)$, consider two open embeddings $U, U' \hookrightarrow Y$ and trivializations $x_U : A|_U \cong U \times \mathbb{C}^{n+1}$ and $x_{U'} : A|_{U'} \cong U' \times \mathbb{C}^{n+1}$, with gluing morphisms over $U \times_Y U'$

$$\varphi = (\varphi_0, \dots, \varphi_n) : (U \times_Y U') \times \mathbb{C}^{n+1} \rightarrow \mathbb{C}^{n+1},$$

such that φ_i is homogeneous of degree w_i with respect to the given weights. For each trivialisation above there is an étale presentation

$$\bigsqcup_{i,j \in \{0, \dots, n\}} (V_{ij} \times U) \rightrightarrows \bigsqcup_{i \in \{0, \dots, n\}} (V_i \times U) \text{ for } P|_{g^{-1}U},$$

$$\text{respectively } \bigsqcup_{i,j \in \{0, \dots, n\}} (V'_{ij} \times U') \rightrightarrows \bigsqcup_{i \in \{0, \dots, n\}} (V'_i \times U') \text{ for } P|_{g^{-1}U'}.$$

Let g_U denote the restriction of g to $P|_{g^{-1}U}$, and similarly for U' . Pullback of the trivialization x_U to $V_i \times_P g_U^*(A|_U)$, of $x_{U'}$ to $V_i \times_P g_U^*(A|_U)$ and of the gluing morphism φ to $((V_i \times U) \times_P (V'_i \times U')) \times \mathbb{C}^{n+1}$ amounts to choosing a root φ_i^{1/w_i} of φ_i such that a point $(V_i \times U) \times_P (V'_i \times U')$ admits a change of coordinates

$$u_i^{j'} = \varphi_j(g(u), p_i(u)) \varphi_i^{-w_j/w_i}(g(u), p_i(u)),$$

where $u = (g(u), (u_i^j)_{j \neq i})$ and $p_i(u) = (u_i^0, \dots, 1, \dots, u_i^n) \in \mathbb{C}^{n+1}$, with 1 in the i -th position. Thus differentiating,

$$\frac{\partial}{\partial u_i^j} = \sum_k \varphi_{kj} \frac{\partial}{\partial u_i^k} (\varphi_i^{-w_k/w_i}) - \varphi_{ij} \left(\sum_k \frac{w_k}{w_i} u_i^k \frac{\partial}{\partial u_i^k} \right) (\varphi_i^{-1}),$$

for $j \neq i$, where the functions $\varphi_{kj} := \frac{\partial \varphi_k}{\partial x_j}$ define a local change of basis for the pullback of the normal bundle $\mathcal{N}_{Y|A}$ and φ_i^{-1/w_i} defines a local change of basis for $\mathcal{O}_P(1)$. Furthermore, as P is a weighted projective bundle, $\varphi_{kj} = 0$ unless $w_k = w_j$

This proves that the trivial extensions of the exact sequence () to $P|_{g^{-1}(U)}$ and $P|_{g^{-1}(U')}$ glue to a restriction of the sequence () on $P|_{g^{-1}(U \times_Y U')}$. $\square$

REFERENCES

- [AGV] Dan Abramovich, Tom Graber, Angelo Vistoli, *Gromov-Witten theory of Deligne-Mumford stacks*, Amer. Journal of Math., Volume 130, Number 5, October 2008, 1337–1398
- [DJS] A. J. de Jong, Jason Michael Starr, *Divisor classes and the virtual canonical bundle for genus 0 maps*, arXiv:math/0602642
- [F] William Fulton, *Intersection Theory*, Springer-Verlag, Berlin, 1984.
- [G] Alexandre Grothendieck, *Séminaire de Géométrie Algébrique du Bois Marie I - 1960-61 - Revêtements tales et groupe fondamental - (SGA 1)* (Lecture notes in mathematics 224). Berlin; New York: Springer-Verlag, xxii+447.

- [K] Andrew Kresch, *Canonical rational equivalence of intersections of divisors*, Invent. Math. 136 (1999) 483–496.
- [L] Valery Lunts, *Coherent Sheaves on Configuration Schemes*, Journal of Algebra, Vol. 244, No.2, 379–406, 2001
- [MM1] A. Mustata, A. Mustata, Intermediate moduli spaces of stable maps, *Invent. Math.* **167** (2007), no. 1, 47–90
- [MM2] A. Mustata, A. Mustata, *The Chow ring of $\overline{M}_{0,m}(n, d)$* , Journal für die Reine und Angewandte Mathematik **615** (2008)
- [10] A. Mustata, A. Mustata, Universal relations on stable map spaces in genus zero, math.AG/0607431
- [P] Rahul Pandharipande, *The canonical class of $\overline{M}_{0,n}(\mathbb{P}^r, d)$ and enumerative geometry*. Internat. Math. Res. Notices 1997, no. 4, 173–186.
- [P] Ian R. Porteous *Blowing up Chern Classes*, Proc. Cambridge Philos. Soc., 56:118–124, 1960.
- [S] *Stack Theory and Applications*. notes taken by H. Clemens at an informal seminar run by A. Bertram, H. Clemens and A. Vistoli; available from www.math.utah.edu/bertram/lectures.
- [V] A. Vistoli, Intersection theory on algebraic stacks and their moduli spaces in Inv. Math. 1989, volume **97**, page 613–670

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY COLLEGE CORK, CORK, IRELAND

MAX PLANCK INSTITUTE FOR MATHEMATICAL SCIENCES

E-mail address: `a.mustata@ucc.ie`, `andrei.mustata@ucc.ie`